



Final

1493 Sixth Line, Oakville

Environmental Impact Study Addendum

Prepared for:

Innovative SHS
117 George Street
Oakville, ON L6J 3B8

Project No. 3096A | August 2025



NATURAL RESOURCE SOLUTIONS INC.

Aquatic, Terrestrial and Wetland Biologists

1493 Sixth Line, Oakville
Environmental Impact Study Addendum

Project Team

| | |
|------------------|---|
| Sydney Gilmour | Project Manager; Terrestrial and Wetland Biologist |
| Ryan Archer | Project Advisor; Senior Terrestrial and Wetland Biologist |
| Ashley Cantwell | Terrestrial and Aquatic Biologist |
| Christy Humphrey | Terrestrial and Wetland Biologist |

Report submitted on August 22, 2025



Sydney Gilmour, M.Sc.
Project Manager
Terrestrial and Wetland Biologist

Table of Contents

| | | |
|------------|---|-----------|
| 1.0 | Introduction | 1 |
| 2.0 | Planning Context..... | 4 |
| 3.0 | Field Methods | 8 |
| 3.1 | Vegetation Community Mapping and Vascular Flora Inventory | 9 |
| 3.2 | Chimney Swift Survey | 9 |
| 3.3 | Bat Habitat and Acoustic Surveys..... | 9 |
| 3.3.1 | Bat Habitat Assessment..... | 9 |
| 3.3.2 | Bat Passive Acoustic Monitoring..... | 10 |
| 3.4 | Headwater Drainage Feature Assessment..... | 11 |
| 3.5 | Wildlife Habitat Assessment and Incidental Wildlife Observations | 13 |
| 4.0 | Biophysical Inventory | 14 |
| 4.1 | Vegetation | 14 |
| 4.1.1 | Vegetation communities..... | 14 |
| 4.1.2 | Vascular Flora | 15 |
| 4.2 | Wildlife..... | 16 |
| 4.2.1 | Birds | 16 |
| 4.2.2 | Herpetofauna..... | 16 |
| 4.2.3 | Mammals | 16 |
| 4.2.4 | Insects | 17 |
| 4.3 | Headwater Drainage Feature..... | 18 |
| 5.0 | Biophysical Analysis..... | 20 |
| 5.1 | Significant Natural Features and Habitats | 20 |
| 5.1.1 | Significant Woodland | 20 |
| 5.1.2 | Headwater Drainage Feature Classification and Management..... | 20 |
| 5.1.3 | Habitat of Threatened and Endangered Species..... | 24 |
| 5.1.4 | Significant Wildlife Habitat | 24 |
| 5.2 | Buffers..... | 24 |
| 6.0 | Impact Assessment, Mitigation and Residual Impacts | 26 |
| 6.1 | Description of the Proposed Work | 26 |
| 6.2 | Approach to Impact Assessment | 26 |
| 6.3 | Direct Impacts | 26 |
| 6.3.1 | Vegetation Removal and Site Grading | 26 |
| 6.3.2 | Impacts to Wildlife and their Habitats | 27 |

| | | |
|-------------|---|-----------|
| 6.4 | Indirect Impacts | 28 |
| 6.4.1 | Disturbance to Adjacent Natural Features and Wildlife Habitat | 28 |
| 6.4.2 | Changes to Hydrological Regime..... | 29 |
| 6.4.3 | Sedimentation and Erosion | 31 |
| 6.4.4 | Water Quality | 32 |
| 6.5 | Induced Impacts | 32 |
| 6.6 | Cumulative Impacts | 33 |
| 6.7 | Residual Impacts | 34 |
| 6.8 | Mitigation..... | 37 |
| 7.0 | Enhancement Opportunities..... | 38 |
| 8.0 | Monitoring Plan | 39 |
| 8.1 | Pre-Construction Monitoring | 39 |
| 8.2 | Construction Monitoring | 39 |
| 8.3 | Post-Construction Monitoring | 39 |
| 9.0 | Conclusion..... | 41 |
| 9.1 | Summary of Recommendations | 42 |
| 10.0 | References | 45 |

List of Tables

| | |
|--|----|
| Table 1. Relevant Policies Legislation and Planning Studies..... | 5 |
| Table 2. Field Survey Summary | 8 |
| Table 3. Acoustic Recorder Settings Used During Bat Passive Acoustic Monitoring (2025). | 11 |
| Table 4. Headwater Drainage Feature Existing Conditions and Management Evaluation | 23 |
| Table 5. Impact Assessment, Mitigation Measures, and Residual Impacts..... | 35 |
| Table 6. Summary of EIS Recommendations..... | 43 |

List of Appendices

Appendix I Proposed Site Plan

Appendix II Terms of Reference

Appendix III Vascular Flora Species Observed within the Subject Property

Appendix IV Bird Species Report from the Study Area

Appendix V Herpetofauna Species Reported from the Study Area

Appendix VI Mammal Species Reported from the Study Area

Appendix VII Lepidoptera Species Reported from the Study Area

Appendix VIII Odonata Species Reported from the Study Area

Maps

Map 1. Study Area

Map 2. Vegetation Communities and Survey Locations

Map 3. Development Constraints

Map 4. Proposed Development Plan

1.0 Introduction

Natural Resource Solutions Inc. (NRSI) was retained by Innovative SHS to complete an Environmental Impact Study (EIS) Addendum associated with proposed residential development located at 1493 Sixth Line, Oakville (hereafter referred to as the subject property; Map 1). In 2024, an EIS was completed on behalf of a previous landowner in support of refinements to the Halton Regional Natural Heritage (RNHS) where it has been mapped on the subject property in the Halton Regional Official Plan (OP) (NRSI 2024). For the purposes of this report, true southwest is referred to as 'west', true northwest is referred to as 'north', etc.

The subject property is approximately 0.87ha in size and contains a single residential dwelling home that was built in 1950, along with two accessory structures (i.e., pigeon shed and lawn shed). The residential dwelling is located within the western section of the property and is surrounded by Cultural Meadow (CUM1), landscaped trees, and a deciduous hedgerow along the northern boundary. A Common Buckthorn (*Rhamnus cathartica*)-dominated Cultural Thicket (CUT1) community is located within the eastern section of the property; a portion of this CUT1 was recently cleared by the landowner. The rear (easternmost) portion of the subject property contains a Cultural Woodland (CUW1) community, which represents a Significant Woodland (NRSI 2024) in the Town of Oakville and Halton Region. The Significant Woodland is also mapped by the Ministry of Natural Resource and Forestry (MNR), which is contiguous with a larger off-site MNR-mapped woodland feature (Map 1), although this connection is fragmented by public trails; the McCraney Valley Trail runs east and north of the subject property. The far east end of the property contains a Headwater Drainage Feature (HDF), which was identified as part the 2024 EIS (NRSI 2024).

The subject property is located within the College Park urban settlement of Oakville. The property is currently zoned as N-Natural Area per the Town of Oakville's Zoning By-law 2014-014 (2025). Prior to 2014, the subject property was zoned and designated as Low Density Residential.

For the purposes of this EIS Addendum, the "study area" is considered the subject property plus adjacent lands within 120m (Map 1). The size of the study area is based on the size of adjacent lands included in the Natural Heritage Reference Manual (OMNR 2010). Adjacent lands include Munn's Public School, commercial businesses, residential neighbourhoods, and recreational fields.

As part of the original EIS (dated May 6, 2024), NRSI carried out a review of background natural heritage information and prepared and submitted a Terms of Reference (TOR) (dated April 23, 2023) to the Region and Town for review and approval. On June 16, 2023, the Region commented that if/when future development is proposed, a new EIS or an EIS Addendum will be required. A new landowner is now proposing to construct a six-storey residential building comprised of 190 rental units on the subject property with an incorporated daycare and amenity space, along with an associated surface-level parking area. See Appendix I for the proposed development plan.

A draft TOR for the EIS Addendum was prepared by NRSI and submitted to the Town of Oakville on May 21, 2025. The Town provided comments on June 23, 2025 and July 15, 2025. A revised TOR (Appendix II) was submitted on July 14, 2025 and July 31, 2025 and approved by the Town on August 1, 2025.

An EIS is required as part of an OP and Zoning Bylaw Amendment application to support the boundary modification of the RNHS designation and zoning on the subject property (see NRSI 2024 for further details). Per Section 16.1.15 a) of Livable Oakville (2009), the specific boundaries of the Natural Area including appropriate buffers of any natural features shall be identified through an EIS.

The EIS must demonstrate that the proposed development (both during and post-construction) will not negatively impact the natural heritage features, including their ecological functions, within the Town and RNHS or unmapped Key Features. Under Section 118.3 of the Regional OP (2024), an Environmental Impact Assessment (otherwise known as an EIS) is required if development or site alteration is proposed within 120m of the RNHS to demonstrate that the proposed development will result in no negative impacts to the RNHS or unmapped Key Features affected by the development by identifying components of the RNHS and assessing the potential environmental impacts, requirements for impact avoidance and mitigation measures, and opportunities for enhancement.

The EIS is required to demonstrate that the proposed OP and Zoning Bylaw Amendment are appropriate for the subject property. As per Section 118.4 of the Regional OP (2024), the recommendations of an EIS are required to be implemented through OP Amendments, Zoning Bylaws, site plan control, conditions of planning approval, or regulations by the appropriate authority

This EIS Addendum summarizes the results of 2025 field surveys and builds on the results of the 2024 EIS (NRSI 2024). The 2023 field survey results, along with the results of the background information review as presented in the 2024 EIS, have been used to characterize the existing natural features on the subject property, including their ecological significance and sensitivity. Recommendations have been provided to avoid, or otherwise effectively mitigate, potential impacts associated with the proposed development. Certain sections from the 2024 EIS (NRSI 2024) have been summarized in this Addendum for completeness and to provide a complete summary of constraints within the subject property. This EIS Addendum was prepared and written in accordance with Halton Region's Environmental Impact Assessment Guidelines (2020).

2.0 Planning Context

For the purposes of this EIS Addendum report, information on the natural heritage features within the study area was collected and assessed for significance. To help inform suitable land-use concepts, guide the layout of development, and identify areas to be protected, inventoried natural features were evaluated against relevant policies, regulations and planning studies (Table 1).

Table 1. Relevant Policies Legislation and Planning Studies

| Policy/Legislation | Description | Project Relevance |
|---|--|--|
| Provincial Planning Statement (PPS; OMMAH 2024). | <ul style="list-style-type: none"> • Issued under the authority of Section 3 of the <i>Planning Act</i> and came into effect on October 20, 2024, replacing the 2020 PPS. • Section 4.1 of the PPS – Natural Heritage establishes clear direction on the adoption of an ecosystem approach and the protection of resources that have been identified as ‘significant’. • Section 4.1.5 of the PPS identifies that development and site alteration shall not be permitted within the area outlined in sub-sections a) – f) “unless it has been demonstrated that there will be no negative impacts on the natural features or their ecological functions.” • The Natural Heritage Reference Manual (OMNR 2010) and the Significant Wildlife Habitat Technical Guide (OMNR 2000, MNRF 2015) were prepared by the MNR to provide guidance on identifying natural features and in interpreting the Natural Heritage sections of the PPS. | <ul style="list-style-type: none"> • Natural features that occur or may occur within the study area, and which receive protection under the PPS, include: <ul style="list-style-type: none"> ○ Significant Woodland, ○ Potential Significant Wildlife Habitat, and ○ Potential habitat for Endangered and Threatened species. |
| <i>Endangered Species Act</i> (2007) | <ul style="list-style-type: none"> • The original ESA, written in 1971, underwent a year-long review which resulted in several changes which came into force in 2007. • The ESA prohibits killing, harming, harassing or capturing Endangered and Threatened species. • The ESA also protects habitat of Endangered and Threatened species from damage and destruction. • Ontario is planning to replace the ESA with the <i>Species Conservation Act</i> (ESA). The exact date the SCA will be in place is yet to be determined by the provincial government. | <ul style="list-style-type: none"> • Based on a preliminary assessment, multiple SAR were identified as having the potential to occur within the study area based on presence of suitable habitat. |
| <i>Protect Ontario by Unleashing our Economy Act, 2025 (Bill 5)</i> | <ul style="list-style-type: none"> • Bill 5, the Protect Ontario by Unleashing our Economy Act, 2025, received Royal Assent on June 5, 2025. • Interim changes to the ESA will be in effect until the SCA is officially in force. Key changes include: • With the passing of Bill 5 on June 5, 2025, habitat is defined as physical dwellings (e.g., dens, nests, hibernacula) and the immediate surrounding area necessary for breeding, rearing, or hibernation. | <ul style="list-style-type: none"> • If SAR protected by the ESA and/or their habitat are confirmed within the study area a mitigation plan, project registration, or permit from the MECP may be required. |

| Policy/Legislation | Description | Project Relevance |
|---|--|--|
| | <ul style="list-style-type: none"> Going forward, proponents will no longer have the option to pay a conservation charge instead of undertaking on-the-ground mitigation. The associated fund will also be dissolved, and the species conservation charge regime will be discontinued. | |
| <i>Migratory Birds Convention Act (MBCA)</i> , 1994 and <i>Migratory Birds Regulations (MBR)</i> , 2022 | <ul style="list-style-type: none"> The MBCA protects migratory game birds, insectivorous birds, and several other migratory non-game birds from persecution in the form of harassment. The schedule of on-site work must consider MBCA timing windows, with timing of breeding bird season typically occurring between April 1 to August 31 (depending on applicable ecozone); however, this is a guideline, since the MBCA applies anytime a migratory bird is nesting. “Incidental take” (unintentional harming, killing, disturbance, or destruction of migratory birds, their nests, or eggs as a result of an activity that is not specifically targeting the bird or nest) is considered illegal, with the exception of a permit obtained by the Canadian Wildlife Service. Schedule 1 of the MBR 2022 provides year-round nest protection for 18 species that are known to re-use nests annually. | <ul style="list-style-type: none"> The timing of construction activities, especially vegetation clearing must have consideration for the MBCA. Typically, this involves avoiding tree and vegetation removal during the peak breeding bird period (April 1 to August 31). |
| <i>Fish and Wildlife Conservation Act (FWCA)</i> ; Government of Ontario 1997) | <ul style="list-style-type: none"> The FWCA provides protection for certain bird species not protected under the MBCA (i.e., raptors), as well as furbearing mammals and their dens or habitual dwellings, aside from the Red Fox (<i>Vulpes vulpes</i>) and Striped Skunk (<i>Mephitis mephitis</i>). | <ul style="list-style-type: none"> The timing of construction activities, especially vegetation clearing and site grading, must have consideration for bird nesting and den sites for fur-bearing mammals. |
| The Livable Oakville Plan: Official Plan (Town of Oakville 2009, last updated April 2025) | <ul style="list-style-type: none"> Land use within the Town of Oakville is guided through the OP that was adopted by the Council of the Corporation of the Town of Oakville on June 22, 2009, and was approved by the Regional Municipality of Halton on November 30, 2009. | <ul style="list-style-type: none"> The subject property is designated Natural Area as per Schedule I Central Land Use. Natural heritage features within the study area, and which receive protection under the Town OP, include: <ul style="list-style-type: none"> Woodlands; Significant Wildlife Habitat; and, Habitat for Endangered and Threatened species. |

| Policy/Legislation | Description | Project Relevance |
|---|--|--|
| | | <ul style="list-style-type: none"> • Per Section 16.1.8, development and site alteration shall not be permitted within regionally significant woodlands or their buffers • Per Section 16.1.10, development and site alteration shall not be permitted in Significant Wildlife Habitat • Per Section 16.1.6, development and site alteration shall not be permitted in the significant habitat of Endangered or Threatened species. Development within 120m of significant habitat of Endangered or Threatened species shall require an EIS to demonstrate that there will be no negative impact on the habitat or its ecological function. |
| Halton Region Official Plan (Region of Halton 2024) | <ul style="list-style-type: none"> • As of July 1, 2025 the Halton Region OP is no longer a regional plan. It is now a Local Plan of the four local municipalities in Halton. • The Halton Region OP identifies the natural features, ecological functions and potential linkages and corridors that comprise the Natural Heritage System (NHS). • The NHS consists of both the Greenbelt Natural Heritage System and the Regional Natural Heritage System. Within the NHS, Key Features are to be protected and maintained for conservation purposes. • Examples of Key Features identified within the Natural Heritage System include significant habitat of SAR, fish habitat, wetlands, ANSI, significant valleylands, significant woodlands, significant wildlife habitat, streams, wetlands, lakes and their littoral zones, seepage areas and springs, aquifers and recharge areas. | <ul style="list-style-type: none"> • The subject property contains the following Key Features: <ul style="list-style-type: none"> • Significant Woodlands; • Significant Wildlife Habitat; and, • Significant habitat of Endangered or Threatened species. • Per Section 139.3.7, it is the policy of the Region to prohibit development or site alteration on lands within Key Features. It is also prohibited to develop or conduct site alteration on lands adjacent to the Key Features unless the proponent has evaluated the ecological functions of these lands through an Environmental Impact Assessment (otherwise known as an EIS). |

3.0 Field Methods

Terrestrial and aquatic field surveys were completed across nine site visits between the period of April – July 2025 to characterize and map the existing natural features present within the subject property. Refer to the 2024 EIS (NRSI 2024) for field survey methodology descriptions for field surveys completed for that report. Table 2 provides a summary of field surveys undertaken within the subject property in 2025.

Table 2. Field Survey Summary

| Date | Field Survey | Protocol | Time | Approximate Person Hours | Weather Conditions | | | | NRSI Staff |
|------------|---|---|-------------|--------------------------|--------------------|--------------------------------|-----------------|-----------------------|---------------------------|
| | | | | | Air Temp. (°C) | Precip. | Cloud Cover (%) | Wind (Beaufort Scale) | |
| 2025-04-15 | Headwater Drainage Feature (HDF) Assessment | Ontario Stream Assessment Protocol (OSAP) (V10.S4.M11) <i>Unconstrained Headwater Sampling</i> module (Gorenc and Stanfield 2017) | 11:00-11:45 | 1.5 | 8 | Rained earlier (light drizzle) | 80 | 2-4 | A. Cantwell C. Kolar |
| 2025-05-08 | | | 15:20-15:40 | 0.75 | 13 | None | 10 | 2-3 | A. Cantwell C. Kolar |
| 2025-07-16 | | | 11:00-11:45 | 0.75 | 8 | Rained earlier | 80 | 2-4 | A. Cantwell |
| 2025-04-22 | Bat Habitat Assessment (Trees and Buildings) | Species at Risk Bats Survey Standard Note - 2022 (MECP 2022a) | 09:30-12:15 | 5.5 | 9 | None | 30 | 3 | C. Humphrey R. Pivar |
| 2025-06-16 | Bat Acoustic Monitoring | Species at Risk Bats Survey Standard Note - 2022 (MECP 2022a), Maternity Roost Surveys (Forests/Woodlands) (MECP 2022b) | - | - | - | - | - | - | C. Humphrey D. Skinner |
| 2025-06-20 | | | - | - | - | - | - | - | A. Van Belleghem |
| 2025-07-03 | | | - | - | - | - | - | - | D. Skinner |
| 2025-07-15 | Vegetation Community Mapping (Ecological Land Classification) | Lee et al. 1998 | 12:00-14:00 | 2 | 30 | None | - | 3 | T. Sieg |
| 2025-07-29 | Chimney Swift Survey | Birds Canada 2023 | 10:29-11:32 | 1 | 26 | None | 9-18 | 2 | M. Alexandrou |

3.1 Vegetation Community Mapping and Vascular Flora Inventory

Vegetation communities were described and mapped according to the Ecological Land Classification (ELC) system for southern Ontario (Lee et al. 1998). This survey included the compilation of a stand description to note the dominant species and cover within each community.

Vegetation community mapping was completed as part of the 2024 EIS (NRSI 2024); however, a section of the Common Buckthorn-dominated Cultural Thicket (CUT1) community had been cleared since the 2024 EIS. Therefore, as part of this EIS Addendum, NRSI reviewed and updated the vegetation community mapping on and adjacent to the subject property. A high-level inventory of vegetation species was also conducted to inform the ELC classifications.

3.2 Chimney Swift Survey

A Chimney Swift (*Chaetura pelagica*) nesting survey was conducted by NRSI staff in accordance with the survey methodology of the *Ontario Swiftwatch Protocol* (Birds Canada 2023) on July 29, 2025. The survey focused on the potential use of the garage turret structure as nesting habitat, since the existing dwelling contains a chimney that is capped and not accessible to breeding Chimney Swifts. The garage turret was surveyed for one hour between 09:00 and one hour before sunset. To improve detectability of the species, the survey was conducted on a day with low cloud cover, low wind, good visibility, and no precipitation.

3.3 Bat Habitat and Acoustic Surveys

3.3.1 Bat Habitat Assessment

Trees

A bat habitat assessment was completed during leaf-off conditions (April 22, 2025) to document potential bat roosting habitat associated with trees (e.g., cracks, crevices, cavities, exfoliating bark, tree species that could provide suitable foliage roosts, etc.) following guidance from the *Species at Risk Bats Survey Note - 2022* (MECP 2022a) and the *Maternity Roost Surveys (Forest/Woodlands)* (MECP 2022b). All trees within the subject property were assessed for potential bat roosting habitat.

All standing live or dead trees with cracks, crevices, hollows, and/or loose or naturally exfoliating bark that could provide suitable roosting habitat for bats, including the SAR Little Brown Myotis

(*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*), and Silver-haired Bat (*Lasionycteris noctivagans*) were documented. Tree species, diameter at breast height (DBH), decay class according to Watt and Caceres (1999), and the number, height, and type (e.g., cavity, crevice, sloughing bark, etc.) of suitable roost sites was documented for each candidate roost tree. The presence of leaf clusters with suitable roosting habitat for Tri-colored Bat (*Perimyotis subflavus*) was also assessed. The density of roost trees was determined for all wooded vegetation communities to characterize habitat availability within the study area.

Structures

Three structures are located within the subject property: a house with an attached garage, a pigeon shed, and a lawn shed. All structures are proposed for removal to accommodate the proposed development. These buildings were assessed for the potential to provide roosting and/or hibernation habitat for bats on April 22, 2025 in accordance with the *Use of Buildings by Species at Risk Bats Survey Methodology* (MECP 2021) and the *Species at Risk Bats Survey Note - 2022* (MECP 2022a).

All external features that may provide suitable roosting habitat or access points to suitable roosting habitat were examined (i.e., fascia, soffits, roofline connections with walls, flashing, siding, etc.). The ground underneath potential access points as well as window sills and walls were also examined during the external inspection for guano and fur oil staining. The structure interiors were also inspected for evidence of potential bat roosting, including attic areas.

Hibernation habitat for SAR bats includes caves, crevices in bedrock extending beyond the frost line, mines containing adits, long concrete culverts, rail tunnels, basements, concrete or stone underground bunkers, and holding tanks with surface accessibility. The basement of the home was examined during interior inspections for potential accessibility to bats and any evidence of bat presence that could indicate hibernation, such as guano.

3.3.2 Bat Passive Acoustic Monitoring

Passive acoustic monitoring was completed in proximity to potential bat roost trees. Bat acoustic monitoring methodology followed the guidelines outlined in the Ministry of Environment, Conservation and Parks (MECP) *Species at Risk Bat Survey Note – 2022* (MECP 2022a), and the MECP's survey protocol for *Maternity Roost Surveys (Forests/Woodlands)* (MECP 2022b).

Two passive acoustic monitoring station locations were selected based on the results of the bat tree habitat assessment. These stations were placed near potential bat roost trees or in suitable foraging and/or travel corridors to assess the potential presence of bats within the subject property (Map 2).

Passive acoustic monitoring was conducted with the use of a Song Meter Mini Bat acoustic recorder (Wildlife Acoustics Inc., Massachusetts, USA) between June 16 and July 3, 2025, for a total of 18 nights, 15 of which were in June. Table 3 summarizes the unit setting used for this project. Data collected on the 10 monitoring nights with the most ideal weather conditions for bat activity (i.e., ambient temperature >10°C, low wind, no precipitation) were selected for further analyses.

Table 3. Acoustic Recorder Settings Used During Bat Passive Acoustic Monitoring (2025).

| Parameter | Setting Used |
|---------------------------|--|
| Detector Type | Wildlife Acoustics Song Meter Mini Bat [full-spectrum] |
| Gain | 12 dB |
| Sample Rate | 384 kHz |
| Minimum Trigger Frequency | 16 kHz |
| Trigger Window | 3 sec |
| Maximum Length | 00:15 min |
| Schedule Start | Sunset + 00:00 hrs |
| Schedule End | Sunset + 05:00 hrs |

Bat echolocation calls recorded during passive acoustic surveys were reviewed with the software program SonoBat 30.2 for the North/Northeastern US & Southern Ontario Region and initially identified to species using the SonoBat Auto-classifier.

Settings for the auto-classification of the acoustic data included the following: auto filter: 5kHz; acceptable call quality: 0.60; decision threshold: 0.90; and maximum number of calls to consider per file: 32. Further manual vetting by NRSI biologists is currently being conducted to confirm species presence and inform potential habitat use type by identified bat species within the subject property.

3.4 Headwater Drainage Feature Assessment

NRSI biologists conducted a headwater drainage feature (HDF) assessment to identify, evaluate and classify HDFs within the subject property. The Evaluation, Classification and

Management of Headwater Drainage Features Guidelines (TRCA and CVC 2014), hereafter referred to as the 'Headwater Guideline', was prepared by the Toronto and Region Conservation Authority (TRCA) and Credit Valley Conservation Authority (CVC) to provide direction for features that are not clearly covered by existing policy and legislation, but may contribute to the overall health and function of the watershed. According to the Headwater Guideline, HDFs include:

- non-permanently flowing drainage features that may not have defined bed or banks;
- first-order and zero-order intermittent and ephemeral channels;
- swales; and
- headwater wetlands.

The Headwater Guideline was developed in conjunction with the Ontario Stream Assessment Protocol (OSAP) Unconstrained Headwater Sampling module (V10.S4.M11) (Gorenc and Stanfield 2017). A feature was identified and investigated within the subject property during a site visit completed on June 9, 2023 with Conservation Halton (CH) staff. The HDF flows in a north to south direction, and drains into a tributary of Sixteen Mile Creek. The HDF falls within the Morrison-Wedgewood Diversion subwatershed catchment area. In 2025, NRSI biologists conducted a full assessment of the HDF in accordance with the methods outlined in the Headwater Guideline and the Ontario Stream Assessment Protocol (OSAP) (V10.S4.M11) Unconstrained Headwater Sampling module (Gorenc and Stanfield 2017). See Map 2 for the HDF reaches.

The feature on the subject property was reclassified from a watercourse to an HDF, prompting the need for a headwater drainage feature assessment in completion of this EIS Addendum. Historically, a culvert or crossing beneath Upper Middle Road directed flow to this feature. However, with the development north of Upper Middle Road, the culvert or crossing was removed, and upstream flows were redirected elsewhere. As a result, the catchment area feeding the feature was significantly reduced, leading to decreased flow in the former watercourse and its reclassification as a headwater drainage feature. During site visits in 2025, the historic channel bed remained visible. See the 2024 EIS (NRSI 2024) for more information.

In the field, each HDF branch was divided into "reaches". These reaches correspond to different sections of the overall feature and were designated based on changes to riparian conditions, channel morphology, and tributary confluences. Each reach was given a unique identifier in the form of "TSMC#-#". Each reach was given the label "TSMC" as the water flows

towards a tributary of Sixteen Mile Creek to the east. Each HDF identified is then assigned a number as they are assessed. Reaches along the same branch are given a secondary number to differentiate them (i.e. TSMC1-1).

For each reach along an HDF, the following data was collected:

- Feature type;
- Flow conditions;
- Flow measurements (if applicable);
- Riparian conditions;
- Feature vegetation;
- Feature and bankfull widths and depths;
- Sediment deposition and transport;
- Site features; and
- Channel connectivity.

NRSI staff conducted three site visits, as per the methods detailed in the Headwater Guideline and the OSAP Unconstrained Headwater Sampling module. The first visit was conducted on April 15, the second on May 8, and the third on July 16, 2025. A full field assessment was conducted for the first site visit which corresponded with spring high water conditions. During the second site visit, the same data was reviewed on site; if any changes occurred since the first site visit, then this additional data was collected. Refinements to reach breaks were also made during the second site visit. The third site visit was to confirm the observations from the first and second site visits and focused on collecting additional hydrology information (i.e., flow conditions and flow measurements, if applicable). The focus of the third site visit was on hydrological conditions and baseflow data collection.

3.5 Wildlife Habitat Assessment and Incidental Wildlife Observations

All incidental observations of wildlife (e.g., mammals, butterflies, odonates (dragonflies and damselflies)) and vegetation species were documented during all field visits. This included both direct and indirect (e.g., tracks, scat, dens, nests, etc.) observations of wildlife presence. Features and species that may be indicative of SWH, as informed by the results of the SWH screening (See the 2024 EIS; NRSI 2024) were documented during the course of all site investigations.

4.0 Biophysical Inventory

4.1 Vegetation

4.1.1 Vegetation communities

Overall, the vegetation communities within the subject property are highly anthropogenically disturbed due to historic use of the land as a residential property and the fragmented nature of the on-site natural features. The natural features within the subject property have been isolated since at least 1985 (Google Earth 2025) due the presence of Munn's Public School to the northeast and open fields to the north and west.

Four vegetation communities were mapped within the subject property, three of which can be classified using the ELC system (Lee et al. 1998). The locations of these communities are shown on Map 2 and the results are described below.

CUW1- Mineral Cultural Woodland Ecosite

As described in the 2024 EIS (NRSI 2024), a cultural woodland community is located at the eastern (rear) extent of the subject property. This feature was categorized as young; there were no indications that the community has undergone a series of natural thinning and replacements (i.e., large diameter trees and down woody debris). The canopy is comprised of Norway Maple (*Acer platanoides*), declining and dead White Ash (*Fraxinus americana*), Manitoba Maple (*Acer negundo*) and White Elm (*Ulmus americana*). The sub-canopy and understorey are not distinguishable due to the age of the feature and comprised an abundance of Common Buckthorn, followed by Dotted Hawthorn (*Crataegus punctata*) and regenerating White Ash. The ground cover consists mainly of common non-native species such as Garlic Mustard (*Alliaria petiolata*), Nipplewort (*Lapsana communis*), Wood Avens (*Geum urbanum*), and Creeping Jennie (*Lysimachia nummularia*).

H- Hedgerow

As described in the 2024 EIS (NRSI 2024) the hedgerow borders the north edge of the cultural meadow and connects to the cultural thicket. The canopy consists of Norway Maple and Siberian Elm (*Ulmus pumila*), while the understorey is dominated by Common Buckthorn. Ground cover consists of occasional Garlic Mustard, Cleavers (*Galium aparine*) and Common Dandelion (*Taraxacum officinale*). The hedgerow is less than 10m wide and less than 0.5ha in size and is not considered an ELC community or a woodland feature. The hedgerow does not

provide any linkage function as it fronts onto Sixth Line and the Munn's Public School parking lot.

CUT1- Mineral Cultural Thicket Ecosite

In 2023, a cultural thicket community previously encompassed the majority of the subject property but the majority of it has since been removed. The cultural thicket provides minimal habitat function, as the dominant species is Common Buckthorn and the ground cover contains greater than 60% Garlic Mustard cover in the spring and summer.

CUM1 – Mineral Cultural Meadow Ecosite

Since the partial removal of the cultural thicket community, a young cultural meadow ecosite has established within the cleared areas and now encompasses the majority of the subject property. The canopy of scattered tree growth consists of Norway Maple (*Acer plantanoides*) and Black Cherry (*Prunus serotina*). The sub-canopy consists of Gray Dogwood species (*Cornus racemosa*), Common Buckthorn and White Ash. The understory is made up of Perennial Sow-Thistle (*Sonchus arvensis*) and Canadian Horseweed (*Erigeron canadensis*). The groundcover is comprised of Black Medic (*Medicago lupulina*) and Garlic Mustard.

4.1.2 Vascular Flora

In total, 84 plants were inventoried within and immediately adjacent to the subject property in 2023 and 2025. A complete list of these species is appended to this report (Appendix III).

No plant SAR or SCC were inventoried within the subject property. No regionally rare species were inventoried within the subject property.

The coefficient of conservatism (CC) is a value ranging from 0 (low) to 10 (high), which is based on a species' tolerance of disturbance and fidelity to a specific habitat integrity (Oldham et al. 1995). Higher values are assigned to species that have specific environmental growing requirements and are less tolerant of disturbance. Average CC value of inventoried plant species on the subject property was 3.14 which is relatively low and indicative of species that are generalist in their habitat preferences and are typically adapted to ecologically disturbed conditions. Of the 84 plant species inventoried, 22 (26%) had CC values of 0-3. Thirty-seven (44%) are non-native in Ontario.

4.2 Wildlife

4.2.1 Birds

In total, 27 bird species were documented within the study area during site investigations in 2023 and 2025. In 2025, six new species were observed within the subject property. Refer to Appendix IV for a list of bird species recorded within in the study area.

One SAR, Chimney Swift, was recorded within the study area. This species is listed as Threatened in Ontario as well as in Canada (MECP 2024, Government of Canada 2024). On July 15, 2025, six Chimney Swift individuals were incidentally observed flying over the subject property. On July 29, 2025, three Chimney Swifts were recorded during a Chimney Swift survey; one was observed flying overhead and two were observed foraging. No Chimney Swifts were observed entering/exiting the garage turret structure, or any on-site structures.

4.2.2 Herpetofauna

NRSI biologists did not observed any herpetofauna during site investigations in 2023 and 2025. A complete list of herpetofauna species reported from the study area and vicinity is provided in Appendix V.

4.2.3 Mammals

In 2025, Northern Raccoon (*Procyon lotor*) (tracks observed; indirect observation) and Eastern Chipmunk (*Tamias striatus*) (direct observation) were recorded within the subject property. In 2023, Eastern Cottontail (*Sylvilagus floridanus*) (direct observation) and Eastern Gray Squirrel (*Sciurus carolinensis*) (direct observation) were recorded within the subject property. All of the mammal species that were recorded are common with secure populations in Ontario (MNR 2024). A complete list of mammals reported from the study area and vicinity is included in Appendix VI.

Bat Habitat Assessment

Trees

Based on the results of the bat habitat assessment, 16 trees were identified as potential roost trees for bats, one of which is located within the dripline of the Cultural Woodland (CUW1). The density of potential roost trees for SAR bats within the CUW1 is 7.7 potential roost trees/ha.

Of those outside of the dripline, 14 were identified as potential roost trees for Little Brown Myotis (*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*) or Silver-haired Bat (*Lasionycteris*

noctivagans) and one tree was identified as a potential roost tree for Tri-colored Bat (*Perimyotis subflavus*). The majority of these trees are located within the hedgerow along the north side of the property.

Structures

House with Attached Garage

The house is a deteriorating wood-sided one and half storey home with an attached garage. The house is still heated (including the basement) and serviced and is occasionally occupied. The interior of the attic space and basement were investigated. While evidence of past use by squirrels was observed, no evidence of use by bats was documented.

Several potential bat entry/exit points were observed in the garage that could provide suitable SAR bat day roosting habitat, and the garage is also easily accessible to mice, squirrels and raccoons (as noted by the presence of scat). However, no bats or bat guano was observed.

Pigeon Shed and Lawn Shed

Both the pigeon shed and lawn shed are dilapidated, with partially collapsed roofs. These buildings were not safe to enter. Given their condition, neither is capable of supporting temperatures required for maternity roosting; however, occasional day-roosting by male or non-reproductive female SAR bats may be possible.

Passive Bat Acoustic Monitoring

Bat acoustic monitoring data is currently undergoing manual vetting of all recorded bat calls that are auto-classified to “rare” SAR bat species (i.e., Northern Myotis, Tri-colored Bat, and Eastern Small-footed Myotis), and a subset of “common” SAR bat species (i.e., Little Brown Myotis, Hoary Bat, Eastern Red Bat, and Silver-haired Bat). Manual vetting will be completed to confirm the recorded call identified to species or species grouping. These results will be summarized in a report and provided under separate cover at a later stage.

4.2.4 Insects

In 2023, one butterfly species, Cabbage White (*Pieris rapae*), was incidentally recorded within the subject property during site investigations. Three additional species, Clouded Sulphur (*Colias philodice*), Black Swallowtail (*Papilio polyxenes*), and Eastern Tiger Swallowtail (*Papilio glaucus*), were incidentally recorded within the subject property in 2025. These species are

relatively common with a secure provincial population (MNR 2024). A complete list of butterfly species reported from the study area vicinity is provided in Appendix VII.

In 2023, one odonate species, Ebony Jewelwing (*Calopteryx maculata*), was incidentally recorded within the subject property during site investigations. This species is relatively common with a secure provincial population (MNR 2024). No odonate species were recorded in 2025. A complete list of odonate species reported from the surrounding 10km squares vicinity is provided in Appendix VIII.

4.3 Headwater Drainage Feature

A single HDF branch, subdivided into two reaches, was determined to be present within the study area. These reaches are described below and shown on Map 2.

Reach TSMC1-1

The TSMC1-1 is 71m long and extends beyond the study area to the southeast. This reach was fully assessed within the subject property upstream of the pedestrian bridge, this portion is a swale that was an historic channel bed. The reach originates at the edge of the Cultural Woodland (CUW1) along the northwest property boundary. The feature flows through the Cultural Woodland community, where it receives input from Reach TSMC1-2 upstream (described below).

During the first HDF visit (April 15, 2025), the reach was minimally flowing. Feature measurements were taken during with the average wetted and feature widths identified as 0.37m and 1.0m, respectively. The average depth of the water was 29mm. During the second visit (May 8, 2025), the reach contained standing water. The average wetted width and depth was 0.48m and 38mm, respectively. The measurements from the second visit are larger due to the targeted areas that still had standing water (pools) within the reach. The entire reach was dry during the third visit (July 16, 2025); however, the substrates were damp throughout the reach (not saturated). The surrounding riparian area was made up of scrubland vegetation, in the form of deciduous shrubs. Substrates were primarily comprised of silt and clay.

An outlet was observed on the eastern bank of the reach at the downstream end. This input was minorly flowing during the first visit. No visible structure was observed; however, it is most likely tile outlet from the soccer fields to the north.

Evidence of sediment transport was observed during the site visits, including erosion on either side of the swale and outlet scour from the presumed tile input from the soccer fields. An average of 70mm of sediment deposition was measured within the feature, classifying it as 'substantial' sediment deposition under the OSAP Unconstrained Headwater Sampling module (Gorenc and Stanfield 2017).

The portion of the reach downstream of the pedestrian bridge is similar to the reach within the subject property. The feature was dry in sections near the second pedestrian crossing (culvert), located approximately 185m downstream of the subject property; this indicates water may be infiltrating. Evidence of historically larger flows were observed as a wide feature bed with poorly formed and eroding banks. As flow inputs upstream were diverted, this feature has developed into a poorly-defined feature.

Reach TSMC1-2

Reach TSMC1-2 is northwest of the subject property. Based on aerial imagery it is an approximately 145m-long grassed swale that conveys flow through a school yard to the TSMC1-1 reach.

Similar to reach TSMC1-1, reach TSMC1-2 was minimally flowing during the first visit, had some pools of standing water during the second visit, and was dry during the third visit. Due to not having site access, measurements were not taken during site visits.

5.0 Biophysical Analysis

An analysis of the significance of existing natural features within the subject property was completed. This analysis is based on the rarity or significance of features and/or associated functions/processes and/or current policies, or planning related studies. Identified significant natural features are described in detail below and are shown on Map 3.

5.1 Significant Natural Features and Habitats

As described above, the study area contains terrestrial and aquatic features and functions that are afforded significance under the Municipal OPs. The following is a summary of the significance and sensitivity of the study area natural features and how the natural heritage policies and legislation inform the identification of constraints for the proposed development.

5.1.1 Significant Woodland

As described in the 2024 EIS (NRSI 2024), the Cultural Woodland (CUW1) community represents a Significant Woodland in Halton Region and is subject to the Region's policies governing this form of Key Feature. Under the Town OP (2009), woodlands are designated as Natural Area lands. In accordance with Town and Region OPs, development or site alteration within or adjacent to (i.e., within 120m of) Regionally Significant Woodlands (as well as their associated buffers) is prohibited unless it has been demonstrated that there will be no negative impacts on the nature feature or its ecological functions (Halton Region 2024, Town of Oakville 2009).

5.1.2 Headwater Drainage Feature Classification and Management

Each HDF reach was assessed to evaluate and classify its functional importance and to identify management recommendations as per the Headwater Guideline (TRCA and CVC 2014). Each reach of the HDF was evaluated based on the four assessment steps outlined in the Headwater Guideline (TRCA and CVC 2014). Step 1 evaluates the hydrologic contribution and function of each reach, Step 2 assesses the riparian vegetation and conditions, Step 3 assesses the feature's contribution to fish and fish habitat, and Step 4 evaluates the terrestrial habitat function each reach provides. The classification results and assessment steps are summarized in Table 4, and the management recommendations for each assessed reach are shown on Map 3

The aquatic habitat assessment in the 2024 EIS (NRSI 2024) was based on one field visit in June 2023 and review of aerial imagery. The feature, which was originally identified as a watercourse, was determined to be an HDF upon further review.

Based on the full three-season field surveys completed in 2025, the overall HDF management recommendation for the reach within the subject property (TSMC1-1) is 'Conservation' and the reach upstream of the subject property (TSMC1-2) is 'Mitigation'. The following management description for 'Conservation' and 'Mitigation' is quoted directly from the Headwater Guideline (TRCA and CVC 2014):

- Conservation – Valued Functions: e.g., seasonal fish habitat with woody riparian cover; marshes with amphibian breeding habitat; or general amphibian habitat with woody riparian cover
 - Maintain, relocate, and/or enhance drainage feature and its riparian zone corridor;
 - If catchment drainage has been previously removed or will be removed due to diversion of stormwater flows, restore lost functions through enhanced lot level controls (i.e., restore original catchment using clean roof drainage), as feasible;
 - Maintain or replace on-site flows using mitigation measures and/or wetland creation, if necessary;
 - Maintain or replace external flows;
 - Use natural channel design techniques to maintain or enhance overall productivity of the reach; and,
 - Drainage feature must connect to downstream.
- Mitigation – Contributing Functions: e.g., contributing fish habitat with meadow vegetation or limited cover
 - Replicate or enhance functions through enhanced lot level conveyance measures, such as well-vegetated swales (herbaceous, shrub and tree material) to mimic online wet vegetation pockets, or replicate through constructed wetland features connected to downstream;
 - Replicate on-site flow and outlet flows at the top end of system to maintain feature functions with vegetated swales, bioswales, etc. If catchment drainage has been previously removed due to diversion of stormwater flows, restore lost functions through enhanced lot level controls (i.e. restore original catchment using clean roof drainage); and,
 - Replicate functions by lot level conveyance measures (e.g. vegetated swales) connected to the natural heritage system, as feasible and/or Low Impact

Development (LID) stormwater options (refer to Conservation Authority Water Management Guidelines for details).

Table 4. Headwater Drainage Feature Existing Conditions and Management Evaluation

| HDF Reach Label | Step 1 | | Step 2 | Step 3 | Step 4 | Management Recommendation |
|--|--|---|---|---|--|---------------------------|
| | Hydrology | Modifier(s) | Riparian Conditions | Fish and Fish Habitat | Terrestrial Habitat | |
| TSMC1-1 | Valued Function Both reaches contained minimally flowing water (<0.5L/s) during the first visit, standing water during the second visit, and both reaches were dry (TSMC1-1 was surface damp) during the third visit. Due to the | A soccer field is adjacent to this feature, tile outlets from the soccer fields may be present. This reach was historically a watercourse, but upstream development within the catchment has reduced its flow. The historic channel bed remained visible during the site visits. | Important Function A forest vegetation community (Cultural Woodland, CUW) is dominant within the riparian corridor of TSMC1-1, resulting in an 'Important' riparian classification. | Contributing Function The entire reach within the subject property was walked during all three visits; no fish were observed in 2025. Barriers to fish exist downstream of the subject property at the pedestrian crossing (culvert). Both reaches function as indirect fish habitat that convey flow, nutrients, and other allochthonous inputs (e.g., invertebrates, organic matter) to aquatic habitats downstream. As such, both reaches have 'Contributing' fish habitat classification. | Limited Function Amphibian habitat is not present within the subject property or upstream. Although, there is cultural woodland there is no habitat upstream of the property. Thus, these reaches do not act like a movement corridor resulting in a 'Limited' function. | CONSERVATION |
| TSMC1-2 (upstream of the subject property) | absence of surface water in July, this reach is categorized as "Valued" hydrology classification. | Evidence of historical and ongoing mowing is present within the grassed feature. | Contributing Function The riparian corridor of TSMC1-2 is dominated by mowed lawn (school yard), resulting in a 'Contributing' riparian classification. | | | MITIGATION |

5.1.3 Habitat of Threatened and Endangered Species

Species at Risk Bats

Treed Bat Habitat

The bat habitat assessment identified 16 trees as potential habitat for SAR bats, 15 of which are outside of the Cultural Woodland (CUW1) community. Fourteen (14) trees are potential habitat for the SAR Little Brown Myotis, Northern Myotis, and/or Silver-haired Bat, and one tree is potential habitat for Tri-colored Bat.

As described in Section 4.2.3, bat passive acoustic monitoring was undertaken to determine if these trees provide bat habitat within the subject property and, if so, identify which species are utilizing this potential habitat. The results of this monitoring are not yet available. Pending the results of that monitoring, the treed features within the subject property, including the 14 potential bat habitat trees, should be considered potential SAR bat habitat.

Anthropogenic Bat Habitat

Based on the habitat assessment, all three structures were identified as capable of supporting day roosting habitat only (i.e., roosting by a male or non-reproductive female, or roosting during migrations). These are relatively low-sensitivity bat habitat functions as individual bats have less fidelity to these specific habitat features and suitable habitat is more widespread on the landscape. In the case of the house, this is limited to the attached garage. The habitat assessment confirmed that these structures do not support more sensitive maternity roosting or hibernation habitat.

5.1.4 Significant Wildlife Habitat

No SWH was confirmed within the study area. Forms of SWH that were initially identified during SWH screening during the TOR stage (Appendix II) were subsequently ruled out based on the results of targeted surveys. See the 2024 EIS (NRSI 2024) for further details.

5.2 Buffers

Buffers are required for natural heritage features to protect them from impacts during and post-construction. At their most basic level, buffers spatially offset development areas from natural features such that direct impacts to the features are avoided. Buffers also represent an important component of a larger suite of recommended measures to mitigate impacts to the adjacent natural features, such as by reducing edge effects. Based on the significance and

sensitivity of the natural features within the study area, ecological buffers must be considered in defining the limits of development on the subject property.

Significant Woodland Buffer

As described in the 2024 EIS (NRSI 2024), woodland buffers are prescribed based on protecting the trees and their root zones as well as providing associated open habitats required by forest wildlife species or for movement. Buffers from woodland driplines are important in maintaining the condition and function of trees within the woodland while protecting them from impacts of adjacent site alteration. A 10m-wide buffer is recommended from the Significant Woodland dripline limit on the subject property (Map 3). The 10m buffer ensures that existing root zones from woodland edge trees will be sufficiently protected while allowing room for future growth, and provides an area of natural woodland edge regeneration and active restoration to enhance the buffering capacity of the feature. A 10m woodland buffer recommendation is in conformance with Section 16.1.8 of the Livable Oakville OP (2009, last updated August 2021),

Headwater Drainage Feature Buffer

Given that the HDF is located within the Significant Woodland, and a 10m buffer is recommended for the Significant Woodland, the HDF will also be buffered.

6.0 Impact Assessment, Mitigation and Residual Impacts

6.1 Description of the Proposed Work

Innovative SHS is proposing the construction of a six-storey residential building comprised of 190 rental units on the subject property with an incorporated daycare and amenity space, along with an associated surface-level parking area. See Map 4 for the proposed development overlaid onto the existing natural features.

6.2 Approach to Impact Assessment

Potential impacts arising from the proposed development are determined by comparing the details of the proposed undertaking with the characteristics of the existing natural features and their functions. The following is a description of the types of impacts that will be discussed, in accordance with the Region's EIA Guidelines (2020).

- Direct impacts to the natural features within the study area associated with disruption or displacement caused by the actual proposed 'footprint' of the development, including impacts caused by site grading and vegetation removal;
- Indirect impacts associated with changes in site conditions such as drainage, water balance and water quantity/quality, and effects of construction on adjacent natural features and habitats;
- Induced impacts associated with continued residential use of the subject property (based on the existing single detached residence), such as disturbance or degradation of the RNHS caused by occupation and use of the property; and,
- Cumulative impacts associated with the spatial and temporal implications of this continued land use in conjunction with other undertakings in the area.

6.3 Direct Impacts

6.3.1 Vegetation Removal and Site Grading

The approach to identifying and delineating the study area's natural features was used to avoid direct impacts from development on significant and sensitive natural features. The proposed development has been designed to avoid direct impacts to the Significant Woodland and HDF. The limits of the proposed development have been set back in accordance with the recommended buffer from these features as shown on Map 4.

The proposed development will be entirely located within the young cultural meadow. No significant vegetation species will be removed as a result of the proposed development.

6.3.2 Impacts to Wildlife and their Habitats

Species at Risk Bat Habitat

The habitat assessment identified 16 trees as potential habitat for SAR bats, 15 of which are outside of the CUW1 woodland that will be retained. Of those outside the woodland, 14 were identified as potential habitat for SAR Little Brown Myotis, Northern Myotis, and/or Silver-haired Bat, and one as potential habitat for Tri-colored Bat. All 14 trees are expected to require removal to accommodate the proposed development. As described in Section 4.2.3, the bat passive acoustic monitoring data are still being analyzed to determine if the on-site treed features are being used by SAR bats; results will be provided at a later stage under separate cover.

Removal of the on-site structures are not anticipated to result in adverse impacts to bat SAR or their habitats, or the contravention of the ESA.

The following mitigation measures are recommended to minimize impacts to bats and their habitats:

- All woody stems >1m in height should be removed outside of the bat active season of March 15-November 30 to avoid direct impacts to individual bats;
- In the event a bat is encountered during construction, work in that area must stop immediately and a qualified biologist consulted to determine next steps; and,
- It is recommended that the limit of construction activities be clearly delineated to avoid encroachment into natural features and habitats.

Other Wildlife Species

Vegetation clearing has the potential to directly impact bird breeding activity through damage and destruction of nests, eggs and young, or avoidance of the area by breeding adults.

Vegetation clearing should therefore occur outside the bird nesting season of April 1-August 31 so as to limit disturbances to nesting activities of birds and to avoid destruction of active nests.

This includes the stripping of herbaceous plant cover from within the agricultural field. The destruction of migratory birds and their nests is prohibited under the federal *Migratory Birds Convention Act*.

6.4 Indirect Impacts

Vegetation clearing, site grading, and construction of the proposed development has the potential to cause indirect impacts to adjacent natural features and functions if not mitigated appropriately. Recommended mitigation measures are provided for each potential impact below.

6.4.1 Disturbance to Adjacent Natural Features and Wildlife Habitat

Vegetation clearing, site grading and other construction activities have the potential to inadvertently destroy, damage and degrade existing vegetation along the development limits unless the development limit boundaries are clearly marked. For example, construction activities can cause scarring and decreased health of adjacent trees whose branches or root systems have been damaged by machinery or affected by construction-related dust and sedimentation. Damage to trees and other vegetation can also be caused by the compaction of soils within tree rooting zones along woodland edges.

Direct damage and indirect disturbances can cause stresses on the natural features that weaken their ecological integrity. In these states, natural features are more prone to establishment and proliferation of invasive, non-native species such as Common Buckthorn, which is already present within the subject property. Proliferation of invasive, non-native species within natural communities decreases their ecological value such as by suppressing native species, diminishing biodiversity and reducing habitat suitability.

To limit ecological impacts during construction, clearly marked construction limits should be established to avoid unnecessary vegetation removal and to ensure that construction activity is maintained outside of these areas. Construction limit fencing should be delineated along the limits of disturbance.

Tree protection fencing must be installed where directed by a Tree Inventory and Preservation Plan (TIPP) and must conform to municipal guidelines in terms of fencing type, signage requirements, etc.

All tree protection fencing must be installed prior to site alteration and construction activities, and inspected by a certified arborist or environmental inspector. Where the need for tree protection and sediment and erosion protection coincides, geotextile materials may be affixed to the bottom of tree protection fencing in accordance with accepted practices. Where tree protection fencing is not required along construction area limits, other forms of boundary

demarcation should be used which may include silt fencing for erosion and sediment control purposes or brightly-coloured snow fencing.

Designated areas for construction lay-down, vehicle access and parking, equipment storage, or materials stockpiling should be located away from the natural features (i.e., Significant Woodland, HDF) and the buffer to limit potential to indirectly impact these features

During construction activities such as vegetation clearing and site grading, dust can potentially result in the following:

- Changes in vegetation due to increased heat absorption and decreased transpiration; and,
- Immediate visual impacts.

Impacts due to dust should be mitigated for by moistening areas of bare, dry soil with water as needed during construction activities to reduce the amount of dust produced.

Excessive noise, vibrations, artificial lighting and human presence as a result of site preparation and construction activities may cause wildlife to temporarily avoid the area. These impacts can be mitigated by restricting the daily timing of construction activities to between 7:00hr and 19:00hr. This timing restriction should also apply to the use of generators or pumps insofar as possible. Any artificial lighting used for construction purposes should be turned off or directed away from the adjacent natural features following the completion of daily construction activities.

Such impacts resulting from dust, noise, and vibrations are expected to be temporary, minimal and localized during the construction of the proposed development. Significant effects on wildlife are not anticipated and it is expected that displaced wildlife species will return to the vicinity of the subject property following construction.

6.4.2 Changes to Hydrological Regime

Surface Water Drainage and Quantity Control

The stormwater management plan for the development has been designed such that catch basin manholes and catch basins will collect drainage and convey flows via private storm sewers to a proposed underground stormwater management tank prior to being discharged into the municipal storm system located on Sixth Line.

Two catchment areas are proposed under post-development conditions: one consisting of paved areas, rooftop areas, and landscaped areas, and another consisting of the wooded area and a concrete walkway (uncontrolled). There will be no additional sheet flow to the rear wooded area (C. Blahut, pers. comm., August 2025). The quantity of runoff will be controlled through an underground storage tank located on the southern section of the subject property as well as a 55m orifice tube to restrict flows exiting the property. The SWM plan for the site will control post-development peak flow rates to pre-development rates between the 2-year and 100-year storm events.

Water Balance

Maintenance of a water balance between pre- and post-development conditions is important to ensure that the hydrological regimes of the receiving aquatic features are not altered through either significant increases or decreases in water inputs. Over the long-term, such imbalances would lead to alterations in the hydrological and ecological functions that these features provide, including but not limited to changes in vegetation community and species composition, and degradation or elimination of certain aquatic and terrestrial habitat functions.

Water balance requires the retention of a 25mm storm event by means of infiltration, evapotranspiration, or reuse. To provide the required volume control to meet site water balance requirements, drainage will be directed to a bottomless tank and a gravel infiltration gallery. Minor drainage will occur to the HDF, however no additional flow will be directed to the feature from pre-development conditions (C. Blahut, pers. comm., August 2025). See Appendix E of the Functional Servicing and Stormwater Management Report (Aplin & Martin Consultants Ltd. 2025) for details.

Interference with Groundwater Flow

Based on a geotechnical study completed by Forward Engineering & Associates Inc. on the subject property (Forward Engineering & Associates Inc. 2025), no groundwater was observed within boreholes, which extended to depths of 1.60-4.67m below the existing ground surface. Due to the depth of groundwater, it is anticipated that subsurface constructions including building foundations/footings and installed servicing infrastructure are unlikely to alter existing groundwater flow patterns. In the event of water seepage into the excavations, it is expected that conventional pumping techniques will be sufficient (Forward Engineering & Associates Inc. 2025).

6.4.3 Sedimentation and Erosion

Construction-Stage

During site stripping and grading activities, areas of bare soil will be exposed which have the potential to erode during rainfall events and impact adjacent natural features such as the rear property woodland and HDF. Increased stormwater surface flow and erosion processes may cause the deposition of sediments onto down-slope vegetation and the adjacent HDF, ultimately causing vegetation die-back or impaired health and a reduction in water quality.

Soil compaction also has potential to occur as a result of heavy machinery in the area of development. Soil compaction can greatly reduce the permeability of soils and affect their ability to retain water during rain/snow melt events. This will result in an increase in surface water run-off which will ultimately increase the erosion potential and the amount of sediment being transported into adjacent natural features.

In order to protect on-site natural features from potential impacts due to sediment, an Erosion and Sediment Control (ESC) Plan must be developed prior to any construction activities on-site. The primary principles associated with sedimentation and erosion protection measures are to: (1) minimize the duration of soil exposure, (2) retain existing vegetation, where feasible, (3) encourage re-vegetation, (4) divert runoff away from exposed soils, (5) keep runoff velocities low, and (6) trap sediment as close to the source as possible.

A detailed ESC Plan should be prepared during the detailed design stage of development planning.

The following general recommendations should be implemented to mitigate erosion and sedimentation impacts, to be refined within the detailed ESC Plan as required:

- Installation of silt fencing along the construction limits in all locations where run-off will discharge to the adjacent natural features. Geotextile material can be attached to tree protection fencing where this fencing type overlaps with silt fencing requirements.
- ESC measures must be regularly inspected and repaired or replaced in a timely manner. Accumulated sediment must be removed as needed.
- Placement of topsoil and seeding of all graded areas not subject to active construction within 30 days. A native seed mix, appropriate to the site conditions and supplemented with a nurse crop, should be applied in areas adjacent to existing natural features.

- It is also recommended that topsoil piles be located away from adjacent natural features and that silt fencing be installed around piles to prevent off-site migration of water-borne sediments.

The impact resulting from soil compaction can be mitigated by minimizing the use of construction vehicles and equipment within buffer areas except where required, and by locating material stockpile and equipment storage locations away from the natural features.

Post-Construction Drainage

As described in Section 6.4.2, post-development site runoff will be controlled to the 1:5-year pre-development levels for storms up to the 1:100-year level. This will be achieved through an underground storage tank and orifice controls.

6.4.4 Water Quality

Decreases in water quality, such as through discharge of deleterious substances in stormwater runoff, can cause both acute and chronic toxicity impacts within biological communities. These impacts include increased mortality rates, impaired health conditions, decreased reproductive productivity and other reproductive impairments in wildlife. Environmental contaminants are also known to biomagnify ‘up the food chain’, where higher-level predators are particularly susceptible to impacts. Water quality impairments can also pose health risks to humans wherever there is potential to come into contact with untreated or inadequately treated water discharge. The water quality of aquatic receptors can also be compromised when excess nutrient concentrations, such as from fertilizers, cause eutrophic conditions which subsequently decrease oxygen availability for fish and other aquatic organisms.

Controlled stormwater runoff will be treated to achieve an “enhanced” level of treatment (80% Total Suspended Solids removal). Stormwater runoff will be conveyed through an Up-Flo Filter. Landscaped areas and rooftops have been deemed inherently clean and have been credited at an 80% removal efficiency (Aplin & Martin Consultants Ltd. 2025). One of the catchment areas will drain to the woodland and a concrete walkway.

6.5 Induced Impacts

Establishment of the proposed development will increase the potential for human disturbances to the adjacent natural features if not properly mitigated. In general, the development may lead

to increased human access to the Significant Woodland and HDF with associated potential for habitat degradation (e.g., vegetation trampling or damage, garbage disposal).

As stated in the 2024 EIS (NRSI 2024), it is recommended that the ecological buffer limit be physically demarcated to ensure that the buffer can be maintained in a natural/restored state and kept outside of actively used portions of the property. This can be achieved by installing permanent fencing along the buffer limit. Installation of permanent fencing along the buffer limit is anticipated to represent an effective deterrence to human encroachment, and the dumping of refuse from the rear of the residential land use, into the natural features and buffer restoration areas.

The proposed development may result in off-site trespassing and garbage dumping/littering by members of the public, particularly due to the presence of an existing pedestrian trail to the immediate rear of the property. Since the Cultural Woodland (CUW1) is part of the greater RNHS, it is not recommended to fence off the rear limits of the property since this may inhibit certain wildlife movements that may occur through the CUW1 community along the wooded corridor that the on-property CUW1 is connected to. It should be noted that there was evidence of human disturbance within the rear portion of the property based on NRSI's site investigations in 2023 and 2025. Therefore, it is recommended that No Trespassing signage be installed adjacent to the current trail to ensure trail users understand that the subject property is private property.

Application of fertilizers and herbicides should not be applied to re-naturalizing vegetation within the buffer. It is recommended that any exterior artificial lighting should be directed away from the adjacent natural features. Exterior lighting fixtures should be downward-casting and Dark Sky-certified (Dark Sky International 2022) to mitigate light pollution effects.

6.6 Cumulative Impacts

In order to evaluate the potential for cumulative impacts resulting from this proposal, it is necessary to look beyond the boundaries of the subject property to the adjacent lands. This approach looks at the character and potential changes that are occurring or may occur in the future on surrounding lands. Cumulative impacts may arise as a result of impacts from a number of sources to add up (or combine) if they overlap in space, overlap in time, occur at some receiver spatially removed from the undertaking, or at some future point in time.

We are not aware of any nearby developments that may affect the subject property natural features (i.e., Significant Woodland and HDF). Provided the recommended mitigation measures presented in this EIS Addendum are implemented, cumulative impacts to the on-site natural features are not anticipated.

6.7 Residual Impacts

Residual impacts to natural features or ecological functions may arise if implemented mitigation measures cannot completely alleviate all impacts. The residual impacts represent the potential effects that may occur, even following implementation of recommended mitigation measures. A summary of potential residual impacts, where they may be expected, is summarized in Table 5.

Table 5. Impact Assessment, Mitigation Measures, and Residual Impacts

| Potential Impact | Development activity creating the impact | Description of impacts by feature and/or function | Mitigation measures | Efficacy and/or residual impacts | Recommended Response |
|---|---|--|--|---|--|
| <ul style="list-style-type: none"> Erosion and sedimentation | <ul style="list-style-type: none"> Site stripping and grading activities Use of heavy machinery in the area of development Stockpiling | <ul style="list-style-type: none"> Exposed areas of bare soil have the potential to erode during rainfall events and cause sediment deposition into adjacent natural features. This is an indirect impact. Soil compaction can reduce the permeability of soils and affect their ability to retain water during rain/snow melts. This is an indirect impact. | <ul style="list-style-type: none"> Prepare and implement a comprehensive Erosion and Sediment Control (ESC) Plan. Heavy-duty ESC fencing is to be installed prior to any vegetation removal, rough grading and construction to demarcate the limit of disturbance. Fencing is to be inspected for proper installation by a qualified inspector and must be maintained for the duration of work until exposed soils stabilize. Any areas of bare soil within the construction area are to be re-vegetated as soon as feasible to prevent erosion of soils and keep dust to a minimum (within 30 days of area being left inactive). An appropriate native seed mix comprised of species is to be applied in areas adjacent to existing natural features Minimize potential for soil compaction. No material stockpile or storage of equipment is to occur within the natural areas. | <ul style="list-style-type: none"> Release of some sediments into natural features. | <ul style="list-style-type: none"> Remove sediment deposits that accumulate outside of the construction limits. Inspect the areas for any signs of residual vegetation damage or disturbance. Review and augment the ESC Plan through additional protective measures where required. |
| <ul style="list-style-type: none"> Damage to/removal of trees and vegetation | <ul style="list-style-type: none"> Vegetation clearing and site grading Use of heavy machinery in the area of development | <ul style="list-style-type: none"> Vegetation clearing and site grading has the potential to inadvertently destroy, damage and degrading existing vegetation along the development limits. Direct damage and indirect disturbances can cause stress on the natural features that weaken their | <ul style="list-style-type: none"> Delineate limits of work zones with heavy-duty ESC fencing. Tree protection fencing must be installed where directed by a Tree Inventory and Preservation Plan (TIPP) and must conform to municipal guidelines. Compensate tree/vegetation removals | <ul style="list-style-type: none"> Death of planted replacement trees, as identified in the TIPP | <ul style="list-style-type: none"> A Certified Arborist or Registered Professional Forester should attend the site as soon as possible to prune damaged tree limbs or roots according to arboricultural best practices. |

| | | | | | |
|---|--|---|---|--|---|
| | | ecological integrity. This may result in the establishment and proliferation of invasive, non-native species. | | | <ul style="list-style-type: none"> All unauthorized materials must be removed from fenced tree protection zones as soon as possible. |
| <ul style="list-style-type: none"> Potential death, injury, or harassment of wildlife | <ul style="list-style-type: none"> Removal of trees and buildings Vegetation clearing Excessive noise, vibrations artificial lighting, and human presence from site preparation (vegetation stripping) and construction activities such as grading. | <ul style="list-style-type: none"> Excessive noise, vibrations, artificial lighting, and human presence due to site preparation and construction activities may cause wildlife to temporarily avoid the area | <ul style="list-style-type: none"> All wood stems >1m in height should be removed outside of the bat SAR active period (i.e., no removals between April 1-November 30). Vegetation clearing should occur outside of the bird nesting season of April 1-August 31. Restrict daily timing of construction activities to between 7:00hr and 19:00hr. Lighting equipment associated with construction activities to be turned off following cessation of daily construction activities, or turned away from natural features. Moisten exposed soils / dry soil with water as needed during construction to reduce dust. | <ul style="list-style-type: none"> No residual impacts are anticipated. | <ul style="list-style-type: none"> N/A |
| <ul style="list-style-type: none"> Human disturbances to the adjacent natural features | <ul style="list-style-type: none"> Establishment of the proposed residential within proximity to the adjacent natural features. | <ul style="list-style-type: none"> The proposed development may lead to increased human access to the adjacent natural features with associated potential for habitat degradation, such as vegetation trampling or damage and garbage disposal. The increase in human access to the natural features is an induced impact. | <ul style="list-style-type: none"> The ecological buffer limit should be physically demarcated to ensure that buffers can be maintained in a natural/restored state. Install No Trespassing signage adjacent to the current trail. | <ul style="list-style-type: none"> No residual impacts are anticipated. | <ul style="list-style-type: none"> N/A |

6.8 Mitigation

The Mitigation Hierarchy is framework that uses an alternatives assessment to minimize negative impacts to the natural heritage system. In accordance with Halton Region's Environmental Impact Assessment Guidelines (2020), mitigation strategies are intended to address or minimize the anticipated and potential impacts such that there is no negative impact resulting from the development or site alteration. The Mitigation Hierarchy typically consists of the following steps:

Step 1: Impact Avoidance

Impact avoidance is the first mitigation measure to consider, as it completely avoids any impacts to natural heritage features.

Step 2: Minimization of Impacts

If impacts cannot be avoided, the next step is to minimize the extent and severity of the impacts. This may include identifying appropriate buffers and developing recommendations for the EIS focused on minimizing impacts such as sediment and erosion control, construction timing windows, etc.

Step 3: Restoration/Enhancement of Existing Natural Features

When impacts cannot be avoided or minimized, then efforts should be made to restore the area to pre-development conditions.

Step 4: Creation of New Natural Features (off-setting)

In some cases, where features are required to be removed to accommodate development, impacts can be mitigated through natural feature creation to address any residual impacts that are not fully addressed by avoidance, minimization, and restoration/enhancement.

The Mitigation Hierarchy was considered in the development planning for the proposed development. The development plan (Map 4) is proposed to be located entirely outside of the natural heritage system (i.e., outside of the HDF and Significant Woodland and its buffer), therefore adhering to Impact Avoidance of the Mitigation Hierarchy.

7.0 Enhancement Opportunities

As recommended in the 2024 EIS (NRSI 2024), the buffer should be planted with a mixture of White Pine (*Pinus strobus*), Eastern White Cedar (*Thuja occidentalis*), Red Oak (*Quercus rubra*), Bur Oak (*Quercus macrocarpa*), Black Cherry (*Prunus serotina*), Sugar Maple (*Acer saccharum*) and native shrubs. In order to suppress buckthorn seedbank regeneration, the conifer component of the planting should account for 50-75% of the planted trees, with the remainder consisting of deciduous species. A native meadow seed mixture, containing species appropriate to the Sixteen Mile Creek subwatershed, should be broadcast throughout the buffer to establish herbaceous cover. The planted species within the buffer should transition from predominantly trees along the eastern edge, closest to the woodland, transitioning to shrubs and meadow edge along the western extent of the buffer.

The restoration plantings would improve the diversity and resiliency of the buffer to mitigate future impacts to the interior woodland area as a result of residential land use (e.g., through attenuation of light casting and noise effects from nearby human occupancy).

The above buffer restoration and enhancement recommendations should be detailed in a future Buffer Restoration and Enhancement Planting Plan and completed to agency satisfaction.

8.0 Monitoring Plan

Pre-, during-, and post-construction monitoring is recommended to ensure that the natural features are not negatively impacted throughout all stages of development. This plan will be implemented through the relevant conditions of planning approvals.

The recommended monitoring components are described below.

8.1 Pre-Construction Monitoring

Prior to any construction activity on-site, including vegetation clearing and site grading, on-site inspections of the following should be undertaken to ensure proper installation:

- Sediment and erosion control measures (e.g., silt fencing) in accordance with an approved ESC Plan; and,
- Tree and natural area protection measures, including proper installation of tree protection fencing in accordance with the Tree Protection Plan.

8.2 Construction Monitoring

Construction monitoring is the responsibility of the proponent and is to be undertaken by a designated environmental inspector or qualified delegate. Generally, construction monitoring must occur to ensure compliance with the conditions of various permits. Construction monitoring measures are to include the following:

- Periodic monitoring of the above measures to ensure maintenance and effectiveness;
- Pruning of any limbs or roots (of trees to be retained) damaged during construction following approved arboricultural techniques;
- Inspection of the ecological buffers to ensure no unauthorized construction encroachments, damage to trees, or other disturbances caused by construction activities outside of the construction limits;
- Fueling and maintenance of machinery to be undertaken at a designated location away from the adjacent natural features and associated buffers; and,
- Storage of machinery and material, fill, etc. in designated areas away from the natural feature and buffer areas.

8.3 Post-Construction Monitoring

It is recommended that the restoration tree and shrub plantings within the Significant Woodland buffer be inspected annually for two years following installation to confirm that the plantings are establishing in good numbers to effectively buffer the woodland. The two-year monitoring window generally coincides with warranty periods for plant materials.

9.0 Conclusion

NRSI was retained by Innovative SHS to complete an EIS Addendum associated with the proposed development of a property located at 1493 Sixth Line in the Town of Oakville. The development would include a six-storey residential building comprised of 190 rental units on the subject property with an incorporated daycare and amenity space, along with an associated surface-level parking area (Map 4). This EIS Addendum was prepared following NRSI's earlier completion of an EIS for a previous owner of the subject property in 2024, which initially characterized the existing on-site natural features. This EIS Addendum builds on the results of the original EIS through completion of additional targeted field surveys.

The subject property contains a Cultural Woodland (CUW1) within the eastern section that represents Significant Woodland within the Town and Region. The on-site portion of Significant Woodland is contiguous with Significant Woodland that extends off and to the rear of the subject property. An HDF was also identified within the eastern section of the property. Based on the full three-season field surveys, the overall HDF management recommendation for the reach within the subject property is 'Conservation' and the reach upstream of the subject property is 'Mitigation'.

A 10m buffer has been recommended from the agency-confirmed Significant Woodland dripline. The 10m Significant Woodland buffer would also spatially offset development areas from the HDF due to its location within the woodland.

Based on the bat habitat assessment, 16 trees were identified as potential habitat for SAR bats, 15 of which are outside of the Significant Woodland. Bat passive acoustic monitoring was undertaken to determine if these trees provide bat habitat within the subject property and, if so, identify which species are utilizing this potential habitat. The results of this monitoring are not yet available. All three on-site structures were identified as capable of supporting day bat SAR roosting habitat only (i.e. roosting by a male or non-reproductive female, or roosting during migrations). In the case of the house, potential bat habitat use is limited to the attached garage. The habitat assessment confirmed that these structures do not support maternity roosting or hibernation habitat. Removal of these structures will not represent a negative impact to SAR bats or their habitat. A Chimney Swift survey was conducted on the subject property; based on these results, ESA-protected habitat for this species was confirmed to be absent on the subject property.

Recommendations were provided to avoid or mitigate impacts to the natural features or their ecological functions. See Section 6.0 for further information on the potential impacts and proposed recommendations.

Based on this study, the proposed development (both during and post-construction) will not negatively impact the natural heritage features, including their ecological functions, within the Town and RNHS or unmapped Key Features, in accordance with Section 118.3 of the Regional OP (2024). Further, the proposed OP and Zoning Bylaw Amendment are appropriate for the subject property.

The bat passive acoustic survey results are still pending; our assessment of SAR bat habitat on the subject property may be updated once completed.

9.1 Summary of Recommendations

The following is a summary of the mitigation measures, enhancement opportunities, and monitoring requirements to be implemented. Please refer to the relevant section of the report for additional details about each recommendation. Where applicable, the summary identifies where these recommendations have been incorporated into existing plans for the development.

Table 6. Summary of EIS Recommendations.

| Recommendations |
|--|
| <i>Direct Impact Mitigation</i> |
| A 10m ecological buffer is recommended from the surveyed Significant Woodland boundary. |
| Vegetation clearing should be maintained outside the period April 1-August 31 to avoid contravention of the <i>Migratory Birds Convention Act</i> . |
| All tree and building removals should occur outside of the active period for the applicable SAR bats (no removals between April 1 - November 30) to avoid direct impacts to individual bats |
| <i>Indirect Impact Mitigation</i> |
| Construction limits must be clearly delineated through use of silt fencing, or other forms of construction fencing. |
| Designated areas for construction lay-down, vehicle access and parking, equipment storage, materials stockpiling and any on-site construction offices should be located away from the natural features and outside of the buffer zone. |
| Dust-prone soils should be moistened with water as needed. |
| The daily timing of construction activities should be maintained to the period 7:00-19:00 hrs. |
| Any artificial lighting used for construction purposes should be turned off or directed away from the adjacent natural features following the completion of daily construction activities. |
| Construction-stage ESC measures must be implemented in accordance with an approved ESC Plan to effectively mitigate erosion and sedimentation impacts on the adjacent natural features. |
| The SWM plan for the site will control post-development peak flow rates to pre-development rates between the 2-year and 100-year storm events. |
| Controlled stormwater runoff will be treated to achieve an “enhanced” level of treatment (80% Total Suspended Solids removal). |
| <i>Induced Impact Mitigations</i> |
| Install No Trespassing signage adjacent to the current trail. |
| The use of pesticides and fertilizers within the buffer area should be avoided. |
| Permanent exterior lighting should be directed away from the woodland, and should be downward-casting to mitigate light pollution effects. |
| Installation of permanent fencing along the buffer to deter human encroachment. |
| <i>Residual Impact Mitigation</i> |
| Remove sediment deposits that accumulate outside of the construction limits. Inspect the areas for any signs of residual vegetation damage or disturbance. Review and augment the ESC Plan through additional protective measures where required. |
| Replacement trees must be inspected two years following their year of planting (coinciding with most nursery stock warranty periods) to ensure their proper establishment and survival. Any replacement trees that are observed to have died or are in poor condition at the time of the inspection should be replaced on a 1:1 basis. |
| <i>Enhancement Opportunities</i> |
| The HDF and Significant Woodland buffer zones should be enhanced with native tree and shrub plantings, as well as a native meadow seed mixture. |
| <i>Monitoring Plan</i> |
| Pre-construction and construction-stage inspections of tree protection and silt fencing are recommended to ensure proper installation and function. |
| Any limbs or roots of trees to be retained that are damaged during construction must be inspected by a certified arborist and pruned where necessary. |

Construction inspections should ensure no unauthorized entry into or damage of buffer and natural features, and that fueling of machinery and stockpiling of materials is maintained away from these areas.

Inspections of enhancement plantings should be undertaken to ensure survival and proper establishment.

10.0 References

- Aplin & Martin Consultants Ltd. 2025. Functional Servicing and Stormwater Management Report. July 10, 2025.
- Birds Canada. 2023. Ontario SwiftWatch: Protocol and Data Forms for Volunteers. Updated March 2023. Birds Canada, Port Rowan, Ontario. Available at: https://www.birds-canada.org/wp-content/uploads/2020/02/Ontario_Swiftwatch_Protocol_2023.pdf
- Forward Engineering & Associates Inc. Geotechnical Investigation. Proposed Multi-Storey Residential Development, 1493 Sixth Line, Oakville, Ontario. June 5, 2025.
- Lee, H.T., W.D. Bakowsky, J. Riley, J. Bowles, M. Puddister, P. Uhlig and S. McMurray. 1998. Ecological Land Classification for Southern Ontario: First Approximation and its Application. Ontario Ministry of Natural Resources, Southcentral Science Section, Science Development and Transfer Branch. SCSS Field Guide FG-02.
- Gorenc, S. and L. Stanfield. 2017. Ontario Stream Assessment Protocol (OSAP). Version 10. Section 4. Module 11. Unconstrained Headwater Sampling.
- Government of Canada. 2024. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2024-12-23. Available: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html>
- Halton Region. 2020. Environmental Impact Assessment Guideline: Regional Official Plan Guidelines.
- Halton Region. 2024. Halton Regional Official Plan. Consolidated May 16, 2024.
- Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>
- Ministry of Environment, Conservation and Parks (MECP). 2022a. Species at Risk Bats Survey Note - 2022. 3pp.
- Ministry of Environment, Conservation and Parks (MECP). 2022b. Maternity Roost Surveys (Forests/Woodlands). 3pp.
- MECP 2021. Ministry of Environment, Conservation and Parks. 2021. Use of Buildings by Species at Risk Bats: Survey Methodology. 3pp
- Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>
- Natural Resource Solutions Inc. (NRSI). 2024. Environmental Impact Study, 1493 Sixth Line, Oakville. May 6, 2024.
- Ontario Ministry of Natural Resources (OMNR). 2010. Natural Heritage Reference Manual for Policies of the Provincial Policy Statement, Second Edition. March 18, 2010.
- Town of Oakville. 202. Town of Oakville Zoning By-law 2014-014. Consolidated 2023.

Town of Oakville. 2009. Livable Oakville: Town of Oakville Official Plan. Last Updated April 22, 2025

Toronto and Region Conservation Authority (TRCA) and Credit Valley Conservation (CVC). 2014. Evaluation, Classification and Management of Headwater Drainage Features Guidelines. January 2014.

Maps



Path: X:\3096_EIS_6thLine_Oakville\NRSI_3096_Map1_StudyArea_2K_2025_08_01_LRC.mxd

Map 1

Sixth Line, Oakville

Study Area

Legend

- Study Area (120m)
- Subject Property
- Headwater Drainage Feature (HDF)
- NRSI Surveyed Dripline Boundary (Reviewed by Halton Region on June 20, 2023)

NATURAL RESOURCE SOLUTIONS INC.
Aquatic, Terrestrial and Wetland Biologists

Map Produced by Natural Resource Solutions Inc. This map is proprietary and confidential and must not be duplicated or distributed by any means without express written permission of NRSI. Data provided by MNR© Copyright: King's Printer Ontario. Imagery: First Base Solutions Inc. (2023).

Project: 3096
Date: August 1, 2025

NAD83 - UTM Zone 17
Size: 11x17"
1:1,750

020406080100

Metres



Map 2

Sixth Line, Oakville

Vegetation Communities and Survey Locations

Legend

- Study Area (120m)
- Subject Property
- Bat Acoustic Survey
- Reach Break
- Headwater Drainage Feature (HDF)
- NRSI Surveyed Dripline Boundary (Reviewed by Halton Region on June 20, 2023)
- Ecological Land Classification (ELC)
- ELC Inclusion
 - (CUM) Cultural Meadow
 - (CUM1) Mineral Cultural Meadow Ecosite
 - (CUT1) Mineral Cultural Thicket Ecosite
 - (CUW1) Mineral Cultural Woodland Ecosite
 - (FOD4) Dry - Fresh Deciduous Forest Ecosite
 - (H) Hedgerow

Map Produced by Natural Resource Solutions Inc. This map is proprietary and confidential and must not be duplicated or distributed by any means without express written permission of NRSI. Data provided by MNR© Copyright: King's Printer Ontario. Imagery: First Base Solutions Inc. (2023).

| | |
|--------------------------------------|--|
| Project: 3096 Date: July 29, 2025 | NAD83 - UTM Zone 17 Size: 11x17" 1:1,700 |
|--------------------------------------|--|

020406080100 Metres

Path: X:\3096_EIS_6thLine_Oakville\NRSI_3096A_Map2_VegetationCommunitiesSurveyLocations_1K_2025_07_29_LRC.mxd



Map 3

Sixth Line, Oakville

Development Constraints

Legend

- Study Area (120m)
- Subject Property
- NRSI Surveyed Dripline Boundary (Reviewed by Halton Region on June 20, 2023)
- Significant Woodland Buffer (10m)
- Reach Break

Headwater Drainage Feature (HDF) Management Recommendation

- Conservation
- Mitigation

Potential Species at Risk Bat Roosting Habitat

- Little Brown Myotis, Northern Myotis, or Silver-haired Bat
- Tri-colored Bat

Ecological Land Classification (ELC)

- ELC
- ELC Inclusion


(CUM) Cultural Meadow
(CUM1) Mineral Cultural Meadow Ecosite
(CUT1) Mineral Cultural Thicket Ecosite
(CUW1) Mineral Cultural Woodland Ecosite
(FOD4) Dry - Fresh Deciduous Forest Ecosite
(H) Hedgerow

Map Produced by Natural Resource Solutions Inc. This map is proprietary and confidential and must not be duplicated or distributed by any means without express written permission of NRSI. Data provided by MNR© Copyright: King's Printer Ontario. Imagery: First Base Solutions Inc. (2023).

Project: 3096
Date: August 1, 2025

NAD83 - UTM Zone 17
Size: 11x17"
1:1,750

Appendix I
Proposed Site Plan



1 SITE PLAN
A-0.1 SCALE: 1:250

RH ZONE REQUIREMENTS

| REQUIREMENT | PROPOSED |
|--|------------------------|
| Minimum lot area* | 1,858.0 m ² |
| Minimum lot frontage | 24.0 m |
| Minimum front yard* | 7.5 m |
| Minimum interior side yard (north) | 4.5 m |
| Minimum interior side yard (south) | 4.5 m |
| Minimum rear yard | 7.5 m |
| Maximum height | n/a |
| Maximum lot coverage* | 35% |
| Minimum landscaping coverage* | 10% |
| Landscape buffer adjacent parking area | 3.0 m |

*based on total lot area (less widening). No deduction for natural area or buffer.

PROPOSED 6-STORY RESIDENTIAL HOUSING

| LOT AREA | GROSS FLOOR AREA |
|---------------------------|------------------------------------|
| GROSS AREA | 8,092.4 m ² (0.8092 ha) |
| ROAD WIDENING | 122.6 m ² (0.0123 ha) |
| LOT AREA: | 7,969.8 m ² (0.7969 ha) |
| NATURAL AREA + 10m BUFFER | 1,803.6 m ² (0.1804 ha) |
| NET LOT AREA: | 6,166.2 m ² (0.6166 ha) |
| 6th FLOOR | 1,852.5 m ² 19,940 sf |
| 5th FLOOR | 1,852.5 m ² 19,940 sf |
| 4th FLOOR | 1,852.5 m ² 19,940 sf |
| 3rd FLOOR | 1,852.5 m ² 19,940 sf |
| 2nd FLOOR | 1,852.5 m ² 19,940 sf |
| 1st FLOOR | 1,852.5 m ² 19,940 sf |
| TOTAL | 11,115.0 m ² 119,640 sf |

UNIT BREAKDOWN

| | |
|-----------|--|
| 6th FLOOR | 27 1br, 7 2br, 1 3br = 35 |
| 5th FLOOR | 27 1br, 7 2br, 1 3br = 35 |
| 4th FLOOR | 27 1br, 7 2br, 1 3br = 35 |
| 3rd FLOOR | 27 1br, 7 2br, 1 3br = 35 |
| 2nd FLOOR | 27 1br, 7 2br, 1 3br = 35 |
| 1st FLOOR | 11 1br, 4 2br, 0 3br = 15 |
| TOTAL | 146 1br, 39 2br, 5 3br = 190 UNITS @ 0.6166 ha = 308 uph |

ACCESSIBLE UNITS

| | |
|-----------|-----------------|
| 1br UNITS | 40 (70%) |
| 2br UNITS | 17 (30%) |
| TOTAL | 57 (30% OF 190) |

RENTABLE AREA


| |
|---------------------------|
| 146 1br @ 415 = 60,590 sf |
| 39 2br @ 600 = 23,400 sf |
| 5 3br @ 850 = 4,250 sf |
| TOTAL = 88,240 sf |

PARKING

| |
|---|
| 190 RENTAL UNITS @ 0.47 = 90 SPACES |
| 450 m ² OFFICE & DAYCARE @ 1/35 = 13 SPACES |
| TOTAL = 103 PARKING SPACES |
| 70 SPACES PROVIDED UNDERGROUND (2,142.0 m ² , 23,055 sf) |
| 33 SPACES ON SURFACE |
| ACCESSIBLE PARKING @ 1, PLUS 3% = 5 REQUIRED, 6 PROVIDED |

FLOOR AREA BREAKDOWN

| | | |
|--|-------------------------|------------|
| FLOOR AREA, RESIDENTIAL | 10,665.0 m ² | 114,815 sf |
| INCLUDES AREA OF ALL SUITES (INCLUDING EXTERIOR WALLS, CORRIDORS, STAIRS, ELEVATORS, LOBBY, ABOVE GRADE SERVICE ROOMS AND RESIDENTIAL AMENITY AREAS) | | |
| FLOOR AREA, NON-RESIDENTIAL | 450.0 m ² | 4,825 sf |
| INCLUDES 1st FLOOR AREAS DEDICATED TO 'HSH' OFFICE, DAYCARE, AND MANAGEMENT OFFICE | | |
| FLOOR AREA, GROSS | 11,115.0 m ² | 119,640 sf |
| ADDITIONAL FLOOR AREAS (NOT INCLUDED IN GROSS AREA) | | |
| BASEMENT PARKING LEVEL | 2,142.0 m ² | 23,055 sf |
| MECHANICAL PENTHOUSE | 82.2 m ² | 885 sf |



pml.A

patrick markus luckie, Architect
7203 BENDIGO CIRCLE - MISSISSAUGA, ON L4M 1Z4 • TEL: 416 855 0100

As drafted, specifications, material quantities and design are the copyright property of Patrick Markus Luckie Architect and must be returned upon request. Reproduction of this drawing, specification, material quantities and design is strictly prohibited without the written permission of Patrick Markus Luckie Architect. This drawing, as an instrument of service, is provided by and is the property of Patrick Markus Luckie Architect and shall remain the property of Patrick Markus Luckie Architect and shall be used only for the project stated on the drawing and shall be returned upon completion of the project. The contractor is responsible for the coordination and verification of all dimensional conditions before any construction and conditions on site as they pertain to these documents. The contractor shall report any discrepancies to the architect in writing prior to the commencement of any official work. Do not make this drawing. This drawing shall not be used for construction purposes unless counterchecked. Patrick Markus Luckie Architect

BLACK DOG
DEVELOPMENT

PENALTA

| Revisions: | |
|------------|-------------|
| Date: | Particular: |
| | |
| | |
| | |
| | |
| | |
| | |

| DATE: | PARTICULAR: |
|---------|------------------------|
| 4.17.25 | ISSUED FOR REVIEW |
| 6.08.25 | ISSUED FOR APPLICATION |
| | |
| | |
| | |

SIXTH LINE
AFFORDABLE HOUSING

1403 SIXTH LINE, OAKVILLE

SITE PLAN

SCALE 1:250

A-0.1

Appendix II
Terms of Reference



NATURAL RESOURCE SOLUTIONS INC.

Aquatic, Terrestrial and Wetland Biologists

July 31, 2025

Project #3096A

To:

Karen Reis, Town of Oakville

Re:

1493 Sixth Line, Oakville

Environmental Impact Study Addendum Terms of Reference

On behalf of Natural Resource Solutions Inc. (NRSI), I am pleased to provide the final Terms of Reference (TOR) for an Environmental Impact Study (EIS) Addendum associated with a proposed residential development on a property located at 1493 Sixth Line, Oakville.

An EIS Addendum will be completed to ensure that the Region of Halton and Town of Oakville natural heritage policies have been addressed.

The attached TOR outlines the steps required to complete the EIS Addendum for the proposed development in accordance with Region and Town requirements.

Should you have any questions or comments regarding this TOR, please do not hesitate to contact the undersigned.

Sincerely,

Natural Resource Solutions Inc.

Sydney Gilmour, M.Sc.

Terrestrial and Wetland Biologist

**1493 Sixth Line, Oakville
Environmental Impact Study Addendum
Final Terms of Reference
July 31, 2025**

Introduction

The subject property is located at 1493 Sixth Line, Town of Oakville, Ontario. The subject property is south of Upper Middle Road East and north of McCraney Street East. See Map 1 for the subject property location. A “study area” has also been identified in order to characterize and assess lands adjacent to (within 120m of) the subject property as site access allows.

An existing residential dwelling is located within the western section of the subject property and is surrounded by manicured grass. The eastern section contains a European Buckthorn (*Rhamnus cathartica*)-dominated Cultural Thicket (CUT1) community; a portion of this feature has recently been cleared by the landowner. The rear (easternmost) portion of the subject property also contains a Cultural Woodland (CUW1) community, which represents a Significant Woodland in the Town of Oakville. A deciduous hedgerow is located along the northern boundary of the subject property adjacent to the residential dwelling and lawns. The far east end of the property contains a Headwater Drainage Feature (HDF). This HDF was identified by NRSI as part of an EIS completed in 2024 on behalf of the previous landowner in support of proposed refinements to the Regional Natural Heritage System (NHS) on the property as mapped in the Halton Region Official Plan (OP) (NRSI 2024). In completion of the 2024 EIS, Conservation Halton confirmed that, as an HDF, the feature is not considered a regulated watercourse.

According to the Regional OP, the entire subject property is designated NHS. Based on this designation, the Town of Oakville’s Zoning By-law has also zoned the subject property as N – Natural Area. The subject property was previously designated and zoned as Low Density Residential, prior to 2014.

It is our understanding that the landowner is proposing to construct a six-storey residential building comprised of 190 rental units on the subject property. The building will also include a daycare, office, bike room, and amenity space, along with associated surface (33) and one level of underground (70) parking spaces.

The purpose of the EIS is to support the boundary modifications of the natural heritage/natural area designation and zoning by-law. An EIS is required to ensure conformance with Regional and Town OP policies, the Provincial Planning Statement (OMMAH 2024), and the *Endangered Species Act*. The purpose of the EIS is to demonstrate that the proposed development will not negatively impact the natural heritage features contained within the Town and Regional NHS or unmapped Key Features and that the proposed OP and zoning amendments are appropriate for the subject property. The EIS will demonstrate that the proposed development (including during- and post-construction) will not have a negative impact on the NHS. The EIS will address Section 118 (3 and 4) of the Region OP and Section 16.1.15 a) and b) of Liveable Oakville.

Characterization of Natural Features

Collection and Review of Background Information

NRSI will utilize background natural heritage and species information that was gathered in completion of the 2024 EIS, including the following resources:

- Ontario Breeding Bird Atlas (BSC et al. 2006);
- Ontario Reptile and Amphibian Atlas (Ontario Nature 2019);
- Atlas of the Mammals of Ontario (Dobbyn 1994); and,
- Ontario Odonata Atlas (OOAD 2023).

The following additional background information sources will be reviewed for updated information to inform this study: on natural heritage features and species records will be collected from the following information sources:

- Natural Heritage Information Centre (NHIC) database of provincially-tracked species (MNR 2025);
- Ontario Butterfly Atlas (Macnaughton et al. 2025); and,
- eBird and iNaturalist online species observations (eBird 2025, iNaturalist 2025).

Background information collection and field survey results completed for the 2024 EIS will serve as a primary source of existing background information in completion of this study.

Species At Risk / Species of Conservation Concern Screening

A screening has been completed to determine the potential for Species at Risk (SAR) and Species of Conservation Concern (SCC) and their habitat to be present within the study area. The habitats within the study area, as determined through completion of the 2024 EIS, have been compared to the habitat requirements of SAR/SCC known from the vicinity of the study area (up to 10km). See Appendix I for the complete SAR/SCC screening table.

Based on the results of the screening, the following SAR that are regulated under the *Endangered Species Act* were identified as having potentially suitable habitat within the study area:

- Silver-haired Bat (*Lasionycteris noctivagans*) – provincially and federally Endangered
- Eastern Red Bat (*Lasiurus borealis*) – provincially and federally Endangered
- Hoary Bat (*Lasiurus cinereus*) – provincially and federally Endangered
- Little Brown Myotis (*Myotis lucifugus*) – provincially and federally Endangered
- Northern Myotis (*Myotis septentrionalis*) – provincially and federally Endangered
- Tri-colored Bat (*Perimyotis subflavus*) – provincially and federally Endangered

Significant Wildlife Habitat Screening

Potential Significant Wildlife Habitat (SWH) types were also screened based previous characterization of the natural features and species habitats within the study area (NRSI 2024) and following discrete significance criteria established by the MNR (MNRF 2015). The results of the SWH screening have informed surveys required to confirm such habitat within the study area, based on any updates to natural feature cover or characteristics within the study area since the 2024 EIS.

Based on the preliminary screening, the following was identified as a Candidate SWH type within the study area, pending further assessment during site investigations:

- Bat Maternity Colonies

See Appendix II for the complete SWH screening tables.

Field Surveys

Field studies have been scoped to characterize and delineate the natural features within the subject property. The following summarizes the field surveys that will be completed to inform the EIS.

Vegetation Community Mapping

NRSI biologists will review and update the vegetation community mapping completed for the 2024 EIS for lands on and adjacent to the subject property. Vegetation community mapping will be completed using the Ecological Land Classification (ELC) system for southern Ontario (Lee *et al.* 1998). A high-level inventory of vegetation species will be conducted (focusing on dominant species) to inform the ELC classifications. Any federally or provincially significant vegetation species that are observed will be documented in detail and GPS-georeferenced.

Headwater Drainage Feature Assessment

NRSI biologists will complete HDF surveys according to the methods outlined in the Headwater Guideline and the Ontario Stream Assessment Protocol (OSAP) (V10.S4.M11) Unconstrained Headwater Sampling module (Gorenc and Stanfield 2017) and the Evaluation, Classification and Management of Headwater Drainage Features Guidelines (CVC and TRCA 2014) to determine the appropriate management. Three site visits will be conducted to capture the early spring high water table conditions (March to mid-April), late spring conditions (late April to mid-May), and summer base-flow conditions (July to August). The HDF will be assessed through four steps to evaluate and classify its functional importance and to identify management recommendations. These steps assess and evaluate the hydrologic contribution and function of the reach, the riparian vegetation and conditions, the feature's contribution to fish, and fish habitat and the terrestrial habitat function the reach provides. The field work included documenting information on ecological and geomorphological form and function to inform these steps.

The full suite of three surveys was not included in the study scope for the 2024 EIS (NRSI 2024), and was recommended to be completed as part of a subsequent EIS on the subject property.

Bat Habitat Assessment

One site visit will be completed to document the presence of any potential bat roosting trees and to assess the existing house (interior and exterior) for evidence of use by bats in accordance with standard protocols (MECP 2021, MECP 2022a, MECP 2022a). This assessment will include the identification of any live trees or snags with tree cavities or loose/sloughing bark that are suitable for roosting. Identified cavity trees will be considered potential habitat for SAR bats. These trees will be recorded in detail on standardized forms and GPS-georeferenced.

Wildlife Habitat Assessment and Incidental Wildlife Observations

Each site visit will include a general assessment of the presence of wildlife habitat within the study area. Any potentially significant habitat will be documented, photographed, and GPS-georeferenced. Observations of all wildlife will be recorded during each site visit, including birds, herpetofauna, mammals, butterflies and odonates (dragonflies and damselflies). In addition to direct observations, any indirect evidence such as dens, tracks, and scat will also be documented.

Natural Feature Constraints Assessment

The results of the field surveys will be combined with any updates to the background information to provide a detailed summary of the existing natural features that occur on and adjacent to the subject property, including any significant habitat features or functions that exist. Potential for significant wildlife species habitat presence will be determined based on updates to the SAR/SCC and SWH screening tables arising from site-level characterization of features and habitat suitability.

All aspects of natural feature significance or sensitivity identified through the background review and site visits will be incorporated into the constraints assessment. An updated constraints map will be prepared for the client, including any recommended refinements to the 10m Significant Woodland buffer proposed in the 2024 EIS (NRSI 2024), to aid in ensuring that the development plan for the lands minimizes or suitably mitigates impacts to the natural features and their ecological functions. This will include the need to avoid or minimize natural feature encroachments.

Impact Assessment, Mitigations, and Recommendations

An impact assessment will be completed based on the details of the proposed development. The assessment will consider potential direct (e.g., habitat removal), indirect (e.g., construction-related, stormwater drainage), and induced (e.g., post-construction human use) impacts on the existing natural features and the ecological functions they provide.

The development plan is proposed to be located outside of the NHS, including the buffer, therefore adhering to the mitigation hierarchy requirements to prioritize opportunities that would avoid impacts to NHS features. The principles of the mitigation hierarchy will be stated and referred to in the EIS, in demonstrating how the plan to avoid direct impacts to the NHS meets this requirement.

Recommendations for key natural heritage feature enhancement and/or restoration will be made where opportunities exist. Recommendations for monitoring will also be provided where applicable, such as to confirm the effectiveness of mitigation measures and to ensure the establishment and survival of enhancement/restoration plantings and seeding.

An EIS will be prepared that includes maps and appendices including taxonomic species lists and a photolog if applicable.

References

- Bird Studies Canada (BSC), Environment Canada's Canadian Wildlife Service, Ontario Nature, Ontario Field Ornithologists and Ontario Ministry of Natural Resources. 2006. Ontario Breeding Bird Atlas Website. Available: <https://www.birdsontario.org/jsp/datasummaries.jsp>
- Credit Valley Conservation (CVC) and Toronto and Region Conservation Authority (TRCA). 2014. Evaluation, Classification and Management of Headwater Drainage Features Guidelines. Finalized January 2014.
- Dobbyn, J.S. 1994. Atlas of the Mammals of Ontario. Don Mills, Federation of Ontario Naturalists. 120p.
- Gorenc, S. and L. Stanfield. 2017. Ontario Stream Assessment Protocol (OSAP). Version 10. Section 4. Module 11. Unconstrained Headwater Sampling.
- Lee, H.T., W.D. Bakowsky, J. Riley, J. Bowles, M. Puddister, P. Uhlig and S. McMurray. 1998. Ecological Land Classification for Southern Ontario: First Approximation and its Application. Ontario Ministry of Natural Resources, Southcentral Science Section, Science Development and Transfer Branch. SCSS Field Guide FG-02.
- Macnaughton A., Layberry R., Cvasin R., Edwards B., and C. Jones. 2025. Ontario Butterfly Atlas. Updated February 2025. Available: <https://www.ontarioinsects.org/atlas/index.html>
- Natural Resource Solutions Inc (NRSI). 2024. Environmental Impact Study (EIS) for 1493 Sixth Line, Oakville. Prepared for Lynda Funk.
- Ontario Ministry of Municipal Affairs and Housing (OMMAH). 2024. Provincial Planning Statement.
- Ontario Ministry of Natural Resources and Forestry (MNRF). 2015. Significant Wildlife Habitat Ecoregion 7E Criterion Schedule: Addendum to Significant Wildlife Habitat Technical Guide. MNRF, January 2015.
- MECP 2022a. Ministry of Environment, Conservation and Parks. 2022a. Species at Risk Bats Survey Note - 2022. 3pp.
- MECP 2022b. Ministry of Environment, Conservation and Parks. 2022b. Maternity Roost Surveys (Forests/Woodlands). 3pp.
- MECP 2021. Ministry of Environment, Conservation and Parks. 2021. Use of Buildings by Species at Risk Bats: Survey Methodology. 3pp.
- Ministry of Natural Resources and Forestry (MNRF). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Ontario Nature. 2019. Reptiles and Amphibians of Ontario Range Maps.
<https://www.ontarioinsects.org/herp/index.html>

Ontario Odonata Atlas Database (OOAD). 2023. Natural Heritage Information Centre, Ontario
Ministry of Natural Resources and Forestry. Species data by 17PJ01 square
accessed on April 22, 2025.

Map



Map 2

Sixth Line, Oakville

Vegetation Communities

Legend

- Subject Property
- Headwater Drainage Feature (HDF)
- NRSI Surveyed Dripline Boundary (Reviewed by Halton Region on June 20, 2023)
- Ecological Land Classification (ELC)
- ELC Inclusion

(CUM) Cultural Meadow

(CUM1) Mineral Cultural Meadow Ecosite

(CUT1) Mineral Cultural Thicket Ecosite

(CUW1) Mineral Cultural Woodland Ecosite

(FOD4) Dry - Fresh Deciduous Forest Ecosite

(H) Hedgerow

NATURAL RESOURCE SOLUTIONS INC.

Aquatic, Terrestrial and Wetland Biologists

Map Produced by Natural Resource Solutions Inc. This map is proprietary and confidential and must not be duplicated or distributed by any means without express written permission of NRSI. Data provided by MNR© Copyright: King's Printer Ontario. Imagery: First Base Solutions Inc. (2023).

| | |
|--------------------------------------|--|
| Project: 3096 Date: July 25, 2025 | NAD83 - UTM Zone 17 Size: 11x17" 1:1,000 |
|--------------------------------------|--|

0102030405060 Metres

Appendix I

Species at Risk and Species of Conservation Concern Habitat Screening

Species at Risk (SAR) and Species of Special Concern (SCC) Screening Table

| Common Name | Scientific Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Habitat Source | Habitat Preference | Suitable Habitats within Study Area | Suitable Habitats within Subject Property | Rationale |
|------------------------|------------------------------|-------|------|---------|------|---------------|--|--|-------------------------------------|---|---|
| Birds | | | | | | | | | | | |
| Northern Bobwhite | <i>Colinus virginianus</i> | S1? | END | E | E | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Grassland, prairie or hay fields with woody cover in form of thickets, tangles of vines, shrubs; fence rows or woodland edges; cropland growing corn, soybeans or small grains and clover or grass; well-drained sandy or loamy soil; pond edges. | No | No | Grassland, prairie, and hay fields are not present within or adjacent to the subject property. |
| Eastern Whip-poor-will | <i>Antrostomus vociferus</i> | S4B | THR | SC | T | Schedule 1 | Recovery Strategy for the Eastern Whip-poor-will (MECP 2019) | Areas with a mix of open and forested areas, such as open woodlands, savannas, pine plantations, woodland edges, or openings in more mature deciduous, coniferous and mixed forests. Forages in open areas and uses forested areas for roosting and nesting. | No | No | Suitable habitat is not present within or adjacent to the subject property. |
| Common Nighthawk | <i>Chordeiles minor</i> | S4B | SC | SC | SC | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Open ground; clearings in dense forests (including burns and logged areas); rock barrens; peat bogs; ploughed fields; gravel beaches or barren areas with rocky soils; open woodlands; flat gravel roofs. | No | No | Undisturbed open ground and forest clearings are not present within or adjacent to the subject property. |
| Chimney Swift | <i>Chaetura pelagica</i> | S3B | THR | T | T | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Commonly found in urban areas near buildings; nests in chimneys, hollow trees, and crevices of rock cliffs. Feeds over open water. | No | No | The on-site structure contains a chimney, however it is capped and not accessible to breeding Chimney Swifts. |
| Eastern Wood-Pewee | <i>Contopus virens</i> | S4B | SC | SC | SC | Schedule 1 | Species at Risk in Ontario (MECP 2024) | Mid-canopy layer of forest clearings and edges of deciduous and mixed forest. Abundant in intermediate-age mature forest stands with little understory vegetation. | No | No | Eastern Wood-Pewee was not recorded during 2023 field surveys. The on-site woodland feature is likely too young to attract breeding Eastern Wood-Pewee. |
| Barn Swallow | <i>Hirundo rustica</i> | S4B | SC | SC | T | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Farmlands, rural areas and other open or semi-open areas near body of water. Nests almost exclusively on human-made structures such as open barns, buildings, bridges and culverts. | No | No | Suitable habitat is not present within or adjacent to the subject property. |
| Purple Martin | <i>Progne subis</i> | S3B | | | | | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Open, trees areas such as farmland, parks, yards, marshes; usually near large bodies of water; colonial; nests in tree cavities, cliff ledges; most common in nest boxes; requires open space for foraging; prefers trees >15 cm dbh. | No | No | Large bodies of water, cavities, and cliff ledges are not present within or adjacent to the subject property. |
| Bank Swallow | <i>Riparia riparia</i> | S4B | THR | T | T | Schedule 1 | Recovery Strategy for the Bank Swallow in Ontario (Falconer et al. 2016) | Nests in burrows in natural and human-made settings with vertical faces in silt and sand deposits. Usually on banks of river and lakes, but also found in sand and gravel pits. | No | No | Banks of rivers and lakes, and sand and gravel pits are not present within or adjacent to the subject property. |
| Tufted Titmouse | <i>Baeolophus bicolor</i> | S3 | | | | | Cornell Lab of Ornithology 2024 | Deciduous woodlands or mixed evergreen-deciduous woodlands with tall trees, typically in areas with a dense canopy and many tree species. Common in orchards, parks, and suburban areas. Generally found at low elevations. | No | No | Woodland does not have a dense canopy layer. Tufted Titmouse was not recorded during 2023 field surveys. |
| Wood Thrush | <i>Hylocichla mustelina</i> | S4B | SC | T | T | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Carolinian and Great Lakes-St. Lawrence forest zones. Undisturbed moist mature deciduous or mixed forest with deciduous sapling growth. Near pond or swamp. Must have some trees higher than 12 m. | No | No | Ponds and swamps are not present within or adjacent to the subject property. |
| Grasshopper Sparrow | <i>Ammodramus savannarum</i> | S4B | SC | SC | SC | Schedule 1 | Significant Wildlife Habitat Technical Guide: Appendix G (OMNR 2000) | Well-drained grassland or prairie with low cover of grasses, taller weeds or sandy soil; hayfields or weedy fallow fields; uplands with ground vegetation of various densities. Requires perches for singing and tracts of grassland generally >5ha. | No | No | Tracts of grassland >5ha in size is not present within or adjacent to the subject property. |

[illegible]

| Common Name | Scientific Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Habitat Source | Habitat Preference | Suitable Habitats within Study Area | Suitable Habitats within Subject Property | Rationale |
|---------------------|----------------------------------|---------|------|---------|------|---------------|---|---|-------------------------------------|---|--|
| Silver-haired Bat | <i>Lasionycteris noctivagans</i> | S3 | END | E | NS | No schedule | COSEWIC Assessment and Status Report on the Hoary Bat <i>Lasiurus cinereus</i> Eastern Red Bat <i>Lasiurus borealis</i> Silver-haired Bat <i>Lasionycteris noctivagans</i> in Canada (COSEWIC 2023) | Roosts primarily in cavities, crevices, and exfoliating bark of typically large-diameter trees in forests and occasionally in or on buildings. Forages in forests of any age, along forests edges and in openings in forests. Primarily overwinters within the U.S. | Possible | Possible | Woodland feature may provide foraging habitat. |
| Eastern Red Bat | <i>Lasiurus borealis</i> | S3 | END | E | NS | No schedule | COSEWIC Assessment and Status Report on the Hoary Bat <i>Lasiurus cinereus</i> Eastern Red Bat <i>Lasiurus borealis</i> Silver-haired Bat <i>Lasionycteris noctivagans</i> in Canada (COSEWIC 2023) | Roosts in foliage of trees; reproductive roosting occurs in upper foliage of typically large-diameter, super-canopy trees in deciduous and coniferous forests of any age. Males occasionally roost in shrubs or saplings. Primarily overwinters within the U.S. | Possible | Possible | Woodland feature may provide suitable roosting habitat. |
| Hoary Bat | <i>Lasiurus cinereus</i> | S3 | END | E | NS | No schedule | COSEWIC Assessment and Status Report on the Hoary Bat <i>Lasiurus cinereus</i> Eastern Red Bat <i>Lasiurus borealis</i> Silver-haired Bat <i>Lasionycteris noctivagans</i> in Canada (COSEWIC 2023) | Roosts in foliage of trees; reproductive roosting occurs in upper foliage of typically large-diameter, super-canopy trees in deciduous and coniferous forests of any age. Forages in the open, such as open wetlands, grasslands and fields with patchy tree cover. Primarily overwinters in the U.S. | Possible | Possible | Woodland feature may provide suitable roosting habitat. |
| Little Brown Myotis | <i>Myotis lucifugus</i> | S3 | END | E | E | Schedule 1 | Recovery Strategy for the Little Brown Myotis, Northern Myotis and Tri-colored Bat in Ontario (Humphrey, C. & H. Fotherby. 2019) | Uses caves, quarries, tunnels, hollow trees or buildings for roosting. Winters in humid caves. Maternity sites in dark warm areas such as attics and barns. Feeds primarily in wetlands and forest edges. | Possible | Possible | The subject property contains buildings that may provide suitable habitat. |
| Northern Myotis | <i>Myotis septentrionalis</i> | S3 | END | E | E | Schedule 1 | Recovery Strategy for the Little Brown Myotis, Northern Myotis and Tri-colored Bat in Ontario (Humphrey, C. & H. Fotherby. 2019) | Roosts in houses and man-made structures but prefers hollow trees or under loose bark. Hibernates in mines or caves. Hunts within forest, below the canopy. | Possible | Possible | The woodland feature may contain suitable roosting trees. The subject property also contain man-made structures. |
| Tri-colored Bat | <i>Perimyotis subflavus</i> | S3? | END | E | E | Schedule 1 | Recovery Strategy for the Little Brown Myotis, Northern Myotis and Tri-colored Bat in Ontario (Humphrey, C. & H. Fotherby. 2019) | Roosts and maternity colonies in older forests and occasionally in barns or other structures. Forage over water and along streams in the forest. Hibernates in caves. | Possible | Possible | The woodland feature may contain suitable roosting trees. |
| Butterflies | | | | | | | | | | | |
| West Virginia White | <i>Pieris virginianensis</i> | S3 | SC | | | | Species at Risk in Ontario (MECP 2024) | Rich, moist, deciduous woods with populations of Two-leaved Toothwort (<i>Cardamine diphylla</i> ; larval food plant). | No | No | Two-leaved Toothwort was not inventoried during the 2023 field surveys. |
| Monarch | <i>Danaus plexippus</i> | S2N,S4B | SC | E | E | Schedule 1 | Species at Risk in Ontario (MECP 2024) | Adults found in a diversity of habitats with a variety of wildflowers. Caterpillars are confined to meadows and open areas where milkweeds grow (larval food plants). | No | No | Milkweed was not inventoried during the 2023 field surveys. |

| Common Name | Scientific Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Habitat Source | Habitat Preference | Suitable Habitats within Study Area | Suitable Habitats within Subject Property | Rationale |
|-----------------|-------------------------------|-------|------|---------|------|---------------|--|---|-------------------------------------|---|--|
| Odonates | | | | | | | | | | | |
| Painted Skimmer | <i>Libellula semifasciata</i> | S3 | | | | | Wisconsin Odonata Survey (2023) | Marshy forest ponds, bogs, slow streams, costal plains. | No | No | Suitable habitat is not present within or adjacent to the subject property. |
| Fish | | | | | | | | | | | |
| Redside Dace | <i>Clinostomus elongatus</i> | S1 | END | E | E | Schedule 1 | Species at Risk in Ontario (MECP 2024) | Pools and slow-moving areas of small streams and headwaters with a gravel bottom. Generally found in areas with overhanging grasses and shrubs. Can be found in shallow parts of streams during spawning. | No | No | A headwater drainage feature is present within eastern section of the subject property, but it does not provide suitable habitat for Redside Dace. |

Reference List

COSEWIC. 2023. COSEWIC assessment and status report on the Hoary Bat *Lasiurus cinereus*, Eastern Red Bat *Lasiurus borealis* and Silver-haired Bat, *Lasionycteris noctivagans*, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xxi + 100 pp. (https://www.cosewic.gc.ca/pdf/publicregistry.html).

Falconer, M., K. Richardson, A. Heagy, D. Tozer, B. Stewart, J. McCracken, and R. Reid. 2016. Recovery Strategy for the Bank Swallow (*Riparia riparia*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. ix + Government of Canada. 2024. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2024-12-23. Available: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html>

Humphrey, C. and H. Fotherby. 2019. Recovery Strategy for the Little Brown Myotis (*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*) and Tri-colored Bat (*Perimyotis subflavus*) in Ontario. Ontario Recovery Strategy Series. Prepared by the Ministry of the Environment, Conservation Linton, J. J. McCarter and H. Fotherby 2018. Recovery Strategy for the Jefferson Salamander (*Ambystoma jeffersonianum*) and Unisexual Ambystoma (Jefferson Salamander dependent population) (*Ambystoma laterale* - (2) *jeffersonianum*) in Ontario. Ontario Recovery Strategy Series.

McCracken, J.D., R.A. Reid, R.B. Renfrew, B. Frei, J.V. Jalava, A. Cowie, and A.R. Couturier. 2013. Recovery Strategy for the Bobolink (*Dolichonyx oryzivorus*) and Eastern Meadowlark (*Stumella magna*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Ministry of the Environment, Conservation and Parks (MECP). 2019. Recovery Strategy for the Eastern Whip-poor-will (*Antrostomus vociferus*) in Ontario. Ontario Recovery Strategy Series. Prepared by the Ministry of the Environment, Conservation and Parks, Peterborough, Ontario. iv + 6 pp. + Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources and Forestry (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Ontario Ministry of Natural Resources (OMNR). 2000. Significant Wildlife Habitat Technical Guide. Appendix G: Wildlife Habitat Matrices and Habitat Descriptions for Rare Vascular Plants. October 2000.

Appendix II
Significant Wildlife Habitat Screening

Significant Wildlife Habitat Assessment Tables

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|--|---|--|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Waterfowl Stopover and Staging Areas (Terrestrial) | | | | | Not Present | Not Present |
| Habitat important to migrating waterfowl | American Black Duck Northern Pintail Gadwall Blue-winged Teal Green-winged Teal American Wigeon Northern Shoveler Tundra Swan | CUM1 CUT1 - Plus evidence of annual spring flooding from melt water or run-off within these Ecosites. - Fields with seasonal flooding and waste grain in the Long Point, Rondeau, Lake. St. Clair, Grand Bend and Pt. Pelee areas may be important to Tundra Swans. | Fields with sheet water during Spring (mid March to May). • Fields flooding during spring melt and run-off provide important invertebrate foraging habitat for migrating waterfowl. • Agricultural fields with waste grains are commonly used by waterfowl, these are not considered SWH unless they have spring sheet water available ^{cxlviii} <u>Information Sources</u> • Anecdotal information from the landowner, adjacent landowners or local naturalist clubs may be good information in determining occurrence. • Reports and other information available from Conservation Authorities (CAs) • Sites documented through waterfowl planning processes (eg. EHJV implementation plan) • Field Naturalist Clubs • Ducks Unlimited Canada • Natural Heritage Information Centre (NHIC) Waterfowl Concentration Area | Studies carried out and verified presence of an annual concentration of any listed species, evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} • Any mixed species aggregations of 100 ^l or more individuals required. • The area of the flooded field ecosite habitat plus a 100-300m radius buffer dependant on local site conditions and adjacent land use is the significant wildlife habitat ^{cxlviii} . • Annual use of habitat is documented from information sources or field studies (annual use can be based on studies or determined by past surveys with species numbers and dates). • SWHMIST ^{cxlix} Index #7 provides development effects and mitigation measures. | Adjacent fields are too small to support aggregations of 100 or more individuals. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Waterfowl Stopover and Staging Areas (Aquatic) | | | | | Not Present | Not Present |
| Important for local and migrant waterfowl populations during the spring or fall migration or both periods combined. Sites identified are usually only one of a few in the eco-district | Canada Goose Cackling Goose Snow Goose Green-winged Teal American Black Duck Northern Pintail Northern Shoveler American Wigeon Gadwall Blue-winged Teal Hooded Merganser Common Merganser Red-breasted Merganser Lesser Scaup Greater Scaup Common Goldeneye Bufflehead Long-tailed Duck Surf Scoter White-winged Scoter Black Scoter Canvasback Redhead Ruddy Duck Brant White-winged Scoter Black Scoter | MAS1 MAS2 MAS3 SAS1 SAM1 SAF1 SWD1 SWD2 SWD3 SWD4 SWD5 SWD6 SWD7 | • Ponds, marshes, lakes, bays, coastal inlets, and watercourses used during migration. Sewage treatment ponds and storm water ponds do not qualify as a SWH, however a reservoir managed as a large wetland or pond/lake does qualify. • These habitats have an abundant food supply (mostly aquatic invertebrates and vegetation in shallow water). <u>Information Sources</u> • Environment Canada • Naturalist clubs often are aware of staging/stopover areas • OMNRF Wetland Evaluations indicate presence of locally and regionally significant waterfowl staging. • Sites documented through waterfowl planning processes (eg. EHJV implementation plan) • Ducks Unlimited projects • Element occurrence specification by Nature Serve: http://www.natureserve.org • Natural Heritage Information Centre (NHIC) Waterfowl Concentration Area | Studies carried out and verified presence of: • Aggregations of 100 ^l or more of listed species for 7 days ^l , results in >700 waterfowl use days. • Areas with annual staging of ruddy ducks, canvasbacks, and redheads are SWH ^{cxlix} • The combined area of the ELC ecosites and a 100m radius area is the SWH ^{cxlviii} • Wetland area and shorelines associated with sites identified within the SWHTG ^{cxlviii} Appendix K ^{cxlix} are significant wildlife habitat. • Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} • Annual Use of Habitat is Documented from Information Sources or Field Studies (Annual can be based on completed studies or determined from past surveys with species numbers and dates recorded). • SWHMIST ^{cxlix} Index #7 provides development effects and mitigation measures. | Ponds, marshes, lakes, bays, coastel inlets, and watercourses are not present within the subject property. Suitable habitat is not present within or adjacent to the subject property. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|---|---|--|--|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Shorebird Migratory Stopover Area | | | | | Not Present | Not Present |
| High quality shorebird stopover habitat is extremely rare and typically has a long history of use | Greater Yellowlegs Lesser Yellowlegs Marbled Godwit Hudsonian Godwit Black-bellied Plover American Golden-Plover Semipalmated Plover Solitary Sandpiper Spotted Sandpiper Semipalmated Sandpiper Pectoral Sandpiper White-rumped Sandpiper Baird's Sandpiper Least Sandpiper Purple Sandpiper Stilt Sandpiper Short-billed Dowitcher Red-necked Phalarope Whimbrel Ruddy Turnstone Sanderling Dunlin | BBO1 BBO2 BBS1 BBS2 BBT1 BBT2 SDO1 SDS2 SDT1 MAM1 MAM2 MAM3 MAM4 MAM5 | Shorelines of lakes, rivers and wetlands, including beach areas, bars and seasonally flooded, muddy and un-vegetated shoreline habitats. Great Lakes coastal shorelines, including groynes and other forms of armour rock lakeshores, are extremely important for migratory shorebirds in May to mid-June and early July to October. Sewage treatment ponds and storm water ponds do not qualify as a SWH. <u>Information Sources</u> • Western hemisphere shorebird reserve network • Canadian Wildlife Service (CWS) Ontario Shorebird Survey • Bird Studies Canada • Ontario Nature • Local birders and naturalist clubs • Natural Heritage Information Center (NHIC) Shorebird Migratory Concentration Area | Studies confirming: • Presence of 3 or more of listed species and > 1000 ^l shorebird use days during spring or fall migration period (shorebird use days are the accumulated number of shorebirds counted per day over the course of the fall or spring migration period). • Whimbrel stop briefly (<24hrs) during spring migration, any site with >100 ^l Whimbrel used for 3 years or more is significant. • The area of significant shorebird habitat includes the mapped ELC shoreline ecosites plus a 100m radius area ^{cxlviii} • Evaluation methods to follow "Bird and Bird Habitats: Guidelines for Wind Power Projects" ^{ccxi} • SWHMIST ^{cxlix} Index #8 provides development effects and mitigation measures. | Lakes, rivers, and wetlands are not present and therefore shoreline habitat is not present. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Raptor Wintering Area | | | | | Not Present | Not Present |
| Sites used by multiple species, a high number of individuals and used annually are most significant | Rough-legged Hawk Red-tailed Hawk Northern Harrier American Kestrel Snowy Owl <u>Special Concern:</u> Short-eared Owl Bald Eagle | <u>Hawks/Owls:</u> Combination of ELC Community Series; need to have present one Community Series from each land class. Forest: FOD, FOM, FOC Upland: CUM, CUT, CUS, CUW <u>Bald Eagle:</u> Forest Community Series: FOD, FOM, FOC, SWD, SWM, or SWC, on shoreline areas adjacent to large rivers or adjacent to lakes with open water (hunting area). | The habitat provides a combination of fields and woodlands that provide roosting, foraging and resting habitats for wintering raptors. Raptor wintering (hawk/owl) sites need to be > 20ha ^{cxlviii, cxlix} with a combination of forest and upland ^{xvi, xvii, xviii, xix, xx, xxi} . Least disturbed sites, idle/fallow or lightly grazed field/meadow (>15ha) with adjacent woodlands ^{cxlix} Field area of the habitat is to be wind swept with limited snow depth or accumulation. Eagle sites have open water and large trees and snags aviable for roosting ^{cxlix} <u>Information Sources</u> • OMNRF Districts • Natural clubs • Natural Heritage Information Centre (NHIC) Raptor Winter Concentration Area • Data from Bird Studies Canada • Reports and other information available from CAs • Results of Christmas Bird Counts | Studies confirm the use of these habitats by: • One or more Short-eared Owls, or, One of more Bald Eagles or; at least 10 individuals and two listed hawk/owl species • To be significant a site must be used regularly (3 in 5 years) ^{cxlix} for a minimum of 20 days by the above number of birds ^l . • The habitat area for an Eagle winter site is the shoreline forest ecosites directly adjacent to the prime hunting area. • Evaluation methods to follow "Bird and Bird Habitats: Guidelines for Wind Power Projects" ^{ccxi} • SWHMIST ^{cxlix} Index #10 and #11 provides development effects and mitigation measures. | Woodlands and adjacent fields are present, but fields are highly disturbed (i.e., recreational fields with human presence) and therefore suitable habitat is not present within or adjacent to the subject property. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|--|--|--|--|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Bat Hibernacula | | | | | Not Present | Not Present |
| Bat hibernacula, are rare habitats in all Ontario landscapes. | Big Brown Bat Eastern Pipistrelle/Tri-colored Bat | Bat Hibernacula may be found in these ecosites: CCR1 CCR2 CCA1 CCA2 (Note: buildings are not considered to be SWH) | Hibernacula may be found in caves, mine shafts, underground foundations and Karsts. Active mine sites should not be considered The locations of bat hibernacula are relatively poorly known. <u>Information Sources</u> • OMNRF for possible locations and contact for local experts • Natural Heritage Information Centre (NHIC) Bat Hibernaculum • Ministry of Northern Development and Mines for location of mine shafts • Clubs that explore caves (eg. Sierra Club) • University Biology Departments with bat experts | • All sites with confirmed hibernating bats are SWH ¹ . • The area includes 200m radius around the entrance of the hibernaculum ^{cxlviii, ccvii, 1} . for the development types and 1000m for wind farms ^{ccv} . • Studies are to be conducted during the peak swarming period (Aug. – Sept.). Surveys should be conducted following methods outlined in the ^{ccv} . "Bats and Bat Habitats: Guidelines for Wind Power Projects" ^{ccv} • SWHMIST ^{cxlix} Index #1 provides development effects and mitigation measures. | Caves, mines shafts, underground foundations, and Karsts are not known to occur in this area. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Bat Maternity Colonies | | | | | Possible | Possible |
| Known locations of forested bat maternity colonies are extremely rare in all Ontario landscapes. | Big Brown Bat Silver-haired Bat | Maternity colonies considered SWH are found in forested Ecosites. All ELC Ecosites in ELC Community Series: FOD FOM SWD SWM | Maternity colonies can be found in tree cavities, vegetation and often in building ^{sxxii, xxv, xxvi, xxvii, xxxi} (buildings are not considered to be SWH). • Maternity roosts are not found in caves and mines in Ontario ^{xxii} . • Maternity colonies located in Mature deciduous or mixed forest stands ^{ccix, ccx} with >10/ha large diameter (>25cm dbh) wildlife trees ^{ccvii} . • Female Bats prefer wildlife tree (snags) in early stages of decay, class 1-3 ^{ccxiv} or class 1 or 2 ^{ccxii} . • Silver-haired Bats prefer older mixed or deciduous forest and form maternity colonies in tree cavities and small hollows. Older forest areas with at least 21 snags/ha are preferred ^{ccx} . <u>Information Sources</u> • OMNRF for possible locations and contact for local experts • University Biology Departments with bat experts | Maternity Colonies with confirmed use by: • >10 Big Brown Bats ¹ • >5 Adult Female Silver-haired Bats ¹ • The area of the habitat includes the entire woodland or the forest stand ELC Ecosite containing the maternity colonies ¹ . • Evaluation methods for maternity colonies should be conducted following methods outlined in the "Bats and Bat Habitats: Guidelines for Wind Power Projects" ^{ccv} . • SWHMIST ^{cxlix} Index #12 provides development effects and mitigation measures. | Suitable tree cavities may be present within the Cultural Woodland (CUW1) feature. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|--|---|--|--|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Turtle Wintering Area | | | | | Not Present | Not Present |
| Generally sites are the only known sites in the area. Sites with the highest number of individuals are most significant. | Midland Painted Turtle <u>Special Concern:</u> Northern Map Turtle Snapping Turtle | Snapping and Midland Painted Turtles: ELC Community Classes: SW, MA, OA and SA ELC Community Series: FEO and BOO Northern Map Turtle: Open Water areas such as deeper rivers or streams and lakes with current can also be used as over-wintering habitat. | <ul style="list-style-type: none">For most turtles, wintering areas are in the same general area as their core habitat. Water has to be deep enough not to freeze and have soft mud substrates.Over-wintering sites are permanent water bodies, large wetlands, and bogs or fens with adequate Dissolved Oxygen^{cix, cx, cxi, cxviii}.Man-made ponds such as sewage lagoons or storm water ponds should not be considered SWH <u>Information Sources</u> <ul style="list-style-type: none">EIS studies carried out by Conservation AuthoritiesField naturalists clubsOMNRF Ecologist or BiologistNatural Heritage Information Centre (NHIC) | <ul style="list-style-type: none">Presence of 5 over-wintering Midland Painted Turtles is significant^l.One or more Northern Map Turtle or Snapping Turtle over-wintering within a wetland is significant^l.The mapped ELC ecosite area with the over wintering turtles is the SWH. If the hibernation site is within a stream or river, the deep-water pool where the turtles are over wintering is the SWH.Over wintering areas may be identified by searching for congregations (Basking Areas) of turtles on warm, sunny days during the fall (Sept. – Oct.) or spring (Mar. – Apr)^{cvii}.Congregation of turtles is more common where wintering areas are limited and therefore significant^{cix, cx, cxi, cxii}.SWHMIST^{cxlix} Index #28 provides development effects and mitigation measures for turtle wintering habitat. | Permanent water bodies, large wetlands, and bogs or fens are not present within or adjacent to the subject property. | |
| Wildlife Habitat: Reptile Hibernaculum | | | | | Not Present | Not Present |
| Generally sites are the only known sites in the area. Sites with the highest number of individuals are most significant | <u>Snakes:</u> Eastern Gartersnake Northern Watersnake Northern Red-bellied Snake Northern Brownsnake Smooth Green Snake Northern Ring-necked Snake <u>Special Concern:</u> Milksnake Eastern Ribbonsnake | For all snakes, habitat may be found in any ecosite in southern Ontario other than very wet ones. Talus, Rock Barren, Crevice and Cave, and Alvar sites may be directly related to these habitats. Observations of congregations of snakes on sunny warm days in the spring or fall is a good indicator. The existence of rock piles or slopes, stone fences, and crumbling foundations assist in identifying candidate SWH. | For snakes, hibernation takes place in sites located below frost lines in burrows, rock crevices and other natural locations. Areas of broken and fissured rock are particularly valuable since they provide access to subterranean sites below the frost line ^{xliv, l, li, lii, cxii} . Wetlands can also be important over-wintering habitat in conifer or shrub swamps and swales, poor fens, or depressions in bedrock terrain with sparse trees or shrubs with sphagnum moss or sedge hummock ground cover. <u>Information Sources</u> <ul style="list-style-type: none">In spring, local residents or landowners may have observed the emergence of snakes on their property (e.g. old dug wells).Reports and other information available from CAsLocal naturalists and experts, as well as university herpetologists may also know where to find some of these sites.Natural Heritage Information Centre (NHIC) | Studies confirming: <ul style="list-style-type: none">Presence of snake hibernacula used by a minimum of five individuals of a snake sp., or, individuals of two or more snake spp.Congregations of a minimum of five individuals of a snake sp., or, individuals of two or more snake spp. near potential hibernacula (eg. foundation or rocky slope) on sunny warm days in Spring (Apr/May) and Fall (Sept/Oct)^l.Note: If there are Special Concern Species present, then site is SWHNote: Sites for hibernation possess specific habitat parameters (e.g. temperature, humidity, etc.) and consequently are used annually, often by many of the same individuals of a local population (i.e. strong hibernation site fidelity). Other critical life processes (e.g. mating) often take place in close proximity to hibernacula. The feature in which the hibernacula is located plus a 30m buffer is the SWH^l.SWHMIST^{cxlix} Index #13 provides development effects and mitigation measures for snake hibernacula. | Rock piles, slopes, stone fences, and crumbling foundaiton are not present within or adjacent to the subject property. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|--|--|---|--|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Colonially - Nesting Bird Breeding Habitat (Bank and Cliff) | | | | | Not Present | Not Present |
| Historical use and number of nests in a colony make this habitat significant. An identified colony can be very important to local populations. All swallow population are declining in Ontario. | Cliff Swallow Northern Rough-winged Swallow (this species is not colonial but can be found in Cliff Swallow colonies) | Eroding banks, sandy hills, borrow pits, steep slopes, and sand piles Cliff faces, bridge abutments, silos, barns Habitat found in the following ecosites: CUM1 CUT1 CUS1 BLO1 BLS1 BLT1 CLO1 CLS1 CLT1 | <ul style="list-style-type: none">Any site or areas with exposed soil banks, undisturbed or naturally eroding that is not a licensed/permitted aggregate area.Does not include man-made structures (bridges or buildings) or recently (2 years) disturbed soil areas, such as berms, embankments, soil or aggregate stockpiles.Does not include a licensed/permitted Mineral Aggregate Operation. <u>Information Sources</u> <ul style="list-style-type: none">Reports and other information available from CAsOntario Breeding Bird Atlas^{ccv}.Bird Studies Canada: Nature Counts http://www.birdscanada.org/birdmon/Field Naturalist clubs | Studies confirming: <ul style="list-style-type: none">Presence of 1 or more nesting sites with 8^{cxlvix} or more cliff swallow pairs and/or rough-winged swallow pairs during the breeding season.A colony identified as SWH will include a 50m radius habitat area from the peripheral nests^{ccvii}.Field surveys to observe and count swallow nests are to be completed during the breeding season. Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects”^{ccxi}.SWHMIST^{cxlix} Index #4 provides development effects and mitigation measures. | Exposed soil banks and associated landscape types are not present within or adjacent to the subject property. | |
| Wildlife Habitat: Colonially - Nesting Bird Breeding Habitat (Tree/Shrubs) | | | | | Not Present | Not Present |
| Large colonies are important to local bird population, typically sites are only known colony in area and are used annually. | Great Blue Heron Black-crowned Night-Heron Great Egret Green Heron | SWM2 SWM3 SWM5 SWM6 SWD1 SWD2 SWD3 SWD4 SWD5 SWD6 SWD7 FET1 | <ul style="list-style-type: none">Nests in live or dead standing trees in wetlands, lakes, islands, and peninsulas. Shrubs and occasionally emergent vegetation may also be used.Most nests in trees are 11 to 15 m from ground, near the top of the tree. <u>Information Sources</u> <ul style="list-style-type: none">Ontario Breeding Bird Atlas^{ccv}, colonial nest records.Ontario Heronry Inventory 1991 available from Bird Studies Canada or NHIC (OMNRF).Natural Heritage Information Centre (NHIC) Mixed Wader Nesting ColonyAerial photographs can help identify large heronries.Reports and other information available from CAsMNRF District OfficesField naturalist clubs | Studies confirming: <ul style="list-style-type: none">Presence of 2 or more active nests of Great Blue Heron or other list species.The habitat extends from the the edge of the colony and a minimum 300m radius or extent of the Forest Ecosite containing the colony or any island <15.0ha with a colony is the SWH^{cc, ccvii}.Confirmation of active colonies must be achieved through site visits conducted during the nesting season (April to August) or by evidence such as the presence of fresh guano, dead young and/or eggshellsSWHMIST^{cxlix} Index #5 provides development effects and mitigation measures. | Wetlands, lakes, islands, and peninsulas are not present within or adjacent to the subject property. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|--|---|---|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Colonially - Nesting Bird Breeding Habitat (Ground) | | | | | Not Present | Not Present |
| Colonies are important to local bird population, typically sites are only known colony in area and are used annually. | Herring Gull Great Black-backed Gull Little Gull Ring-billed Gull Common Tern Caspian Tern Brewer's Blackbird | Any rocky island or peninsula (natural or artificial) within a lake or large river (two-lined on a 1:50,000 NTS map). Close proximity to watercourses in open fields or pastures with scattered trees or shrubs (Brewer's Blackbird) MAM1 – 6 MAS1 – 3 CUM CUT CUS | <ul style="list-style-type: none">• Nesting colonies of gulls and terns are on islands or peninsulas associated with open water or in marshy areas.• Brewers Blackbird colonies are found loosely on the ground in or in low bushes in close proximity to streams and irrigation ditches within farmlands. <u>Information Sources</u> <ul style="list-style-type: none">• Ontario Breeding Bird Atlas^{ccv}, rare/colonial species records.• Canadian Wildlife Service• Reports and other information available from CAs• Natural Heritage Information Centre (NHIC) Colonial Waterbird Nesting Area• MNRF District Offices• Field naturalist clubs | Studies confirming: <ul style="list-style-type: none">• Presence of >25 active nests for Herring Gulls, >5 active nests for Common Tern or >2 active nests for Caspian Tern^l.• Any active nesting colony of one or more Little Gull, and Great Black-backed Gull is significant^l.• Presence of 5 or more pairs for Brewer's Blackbird^l.• The edge of the colony and a minimum 150m radius area of the habitat, or the extent of the ELC ecosites containing the colony or any island <3.0ha with a colony is the SWH^{cc, ccvii}.• Studies would be done during May/June when actively nesting. Evaluation methods to follow "Bird and Bird Habitats: Guidelines for Wind Power Projects"^{ccxi}.• SWHMIST^{cxlix} Index #6 provides development effects and mitigation measures. | Rocky islands or peninsulas within a lake or large river are not located within or adjacent to the subject property. | |
| Wildlife Habitat: Migratory Butterfly Stopover Areas | | | | | Not Present | Not Present |
| <u>Rationale:</u> Butterfly stopover areas are extremely rare habitats and are biologically important for butterfly species that migrate south for the winter | Painted Lady Red Admiral <u>Special Concern:</u> Monarch | Combination of ELC Community Series; need to have present one Community Series from each landclass: Field: CUM CUT CUS Forest: FOC FOD FOM CUP Anecdotally, a candidate site for butterfly stopover will have a history of butterflies being observed. | A butterfly stopover area will be a minimum of 10ha in size with a combination of field and forest habitat present, and will be located within 5km of Lake Ontario and Erie ^{cxlix} . <ul style="list-style-type: none">• The habitat is typically a combination of field and forest, and provides the butterflies with a location to rest prior to their long migration south^{xxxii, xxxiii, xxxiv, xxxv, xxxvi}.• The habitat should not be disturbed, fields/meadows with an abundance of preferred nectar plants and woodland edge providing shelter are requirements for this habitat^{cxlviii, cxlix}.• Staging areas usually provide protection from the elements and are often spits of land or areas with the shortest distance to cross the Great Lakes^{xxxvii, xxxviii, xxxix, xl, xli}. <u>Information Sources</u> <ul style="list-style-type: none">• MNRF District Offices• Natural Heritage Information Centre (NHIC)• Agriculture Canada in Ottawa may have list of butterfly experts.• Field Naturalist Clubs• Toronto Entomologists Association• Conservation Authorities | Studies confirm: <ul style="list-style-type: none">• The presence of Monarch Use Days (MUD) during fall migration (Aug/Oct)^{xliii}. MUD is based on the number of days a site is used by Monarchs, multiplied by the number of individuals using the site. Numbers of butterflies can range from 100-500/day^{xxxvii}, significant variation can occur between years and multiple years of sampling should occur^{xl, xlii}.• Observational studies are to be completed and need to be done frequently during the migration period to estimate MUD• MUD of >5000 or >3000 with the presence of Painted Ladies or White Admiral's is to be considered significant^l.• SWHMIST^{cxlix} Index #16 provides development effects and mitigation measures. | The subject property is located within 5km of Lake Ontario and forest and field habitat are present. However, the study area is highly disturbed; adjacent lands include recreational fields, schools, and residential areas. | |

Table 1. Characteristics of Seasonal Concentration Areas for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|--|--|--|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Landbird Migratory Stopover Areas | | | | | Not Present | Not Present |
| Sites with a high diversity of species as well as high numbers are most significant | All migratory songbirds Canadian Wildlife Service Ontario website: http://www.on.ec.gc.ca/wildlife_e.html All migrant raptors species Ontario Ministry of Natural Resources: Fish and Wildlife Conservation Act, 1997. Schedule 7: Specially Protected Birds (Raptors) | All Ecosites associated with these ELC Community Series: FOC FOM FOD SWC SWM SWD | Woodlots need to be >5 ha ¹ in size and within 5km ^{iv, v, vi, vii, viii, ix, x, xi, xii, xiii, xiv, xv} of Lake Ontario and Erie. If woodlands are rare in an area of shoreline, woodland fragments 2-5ha can be considered for this habitat • If multiple woodlands are located along the shoreline those Woodlands <2km from Lake Erie or Ontario are more significant ^{cxlix} . • Sites have a variety of habitats: forest, grassland and wetland complexes ^{cxlix} . • The largest sites are more significant ^{cxlix} • Woodlots and forest fragments are important habitats to migrating birds ^{ccxviii} , these features located along the shore and located within 5km of Lake Ontario and Lake Erie are Candidate SWH ^{cxlviii} . <u>Information Sources</u> • Bird Studies Canada • Ontario Nature • Local birders and naturalist clubs • Ontario Important Bird Areas (IBA) Program | Studies confirm: • Use of the habitat by >200 birds/day and with >35 spp. with at least 10 bird spp. recorded on at least 5 different survey dates ¹ . This abundance and diversity of migrant bird species is considered above average and significant. • Studies should be completed during spring (March/May) and fall (Aug/Oct) migration using standardized assessment techniques. Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} . • SWHMIST ^{cxlix} Index #9 provides development effects and mitigation measures. | The subject property is located within 5km of Lake Ontario, and the woodland is larger than 5ha in size. However, abundance (>200 birds/day) and diversity (>35 spp.) were not observed during 2023 field surveys. The wooded community within the subject property was identified as CUW1 and therefore does not meet the ELC ecosite criteria to support landbird migratory stopover areas. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Deer Winter Congregation Areas | | | | | Not Present | Not Present |
| Deer movement during winter in the southern areas of Ecoregion 7E are not constrained by snow depth, however deer will annually congregate in large numbers in suitable woodlands to reduce or avoid the impacts of winter conditions ^{cxlviii} | White-tailed Deer | All Forested Ecosites with these ELC Community Series: FOC FOM FOD SWC SWM SWD Conifer plantations (CUP) smaller than 50 ha may also be used. | • Woodlots >100 ha in size or if large woodlots are rare in a planning area woodlots>50ha ¹ . • Deer movement during winter in Ecoregion 7E are not constrained by snow depth, however deer will annually congregate in large numbers in suitable woodlands ^{cxlviii} . • Large woodlots > 100ha and up to 1500 ha are known to be used annually by densities of deer that range from 0.1-1.5 deer/ha ^{ccxxiv} . • Woodlots with high densities of deer due to artificial feeding are not significant ¹ . <u>Information Sources</u> • MNRF District Offices • LIO/NRVIS | Studies confirm: • Deer management is an MNRF responsibility, deer winter congregation areas considered significant will be mapped by MNRF ^{cxlviii} . • Use of the woodlot by white-tailed deer will be determined by MNRF, all woodlots exceeding the area criteria are significant, unless determined not to be significant by MNRF ¹ . • Studies should be completed during winter (Jan/Feb) when >20cm of snow is on the ground using aerial survey techniques ^{ccxxiv} , ground or road surveys, or a pellet count deer density survey ^{ccxxv} . • SWHMIST ^{cxlix} Index #2 provides development effects and mitigation measures. | Deer wintering area is not mapped by the Ministry of Natural Resources. Further, the woodland does not meet the >100ha size criterion for this SWH. Suitable habitat is not present within or adjacent to the subject property. | |

Significant Wildlife Habitat Assessment Tables

Table 2. Characteristics of Rare Vegetation Communities for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|--|--|--|---|------------------|
| | ELC Ecosite Codes | Habitat Description | Detailed Information and Sources | Defining Criteria | Study Area | Subject Property |
| Cliff and Talus Slopes | | | | | Not Present | Not Present |
| Cliffs and Talus Slopes are extremely rare habitats in Ontario. | Any ELC Ecosite within Community Series: TAO TAS TAT CLO CLS CLT | A Cliff is vertical to near vertical bedrock >3m in height. A Talus Slope is rock rubble at the base of a cliff made up of coarse rocky debris. | Most cliff and talus slopes occur along the Niagara Escarpment. <u>Information Sources</u> <ul style="list-style-type: none">• The Niagara Escarpment Commission has detailed information on location of these habitats.• OMNRF Districts• Natural Heritage Information Centre (NHIC) has location information available on their website• Field naturalist clubs• Conservation Authorities | <ul style="list-style-type: none">• Confirm any ELC Vegetation Type for Cliffs or Talus Slopes^{lxxviii}• SWHMIST^{cxlix} Index #21 provides development effects and mitigation measures. | Cliff and Talus Slopes are not present within or adjacent to the subject property | |
| Sand Barrens | | | | | Not Present | Not Present |
| Sand barrens are rare in Ontario and support rare species. Most Sand Barrens have been lost due to cottage development and forestry. | ELC Ecosites: SBO1 SBS1 SBT1 Vegetation cover varies from patchy and barren to continuous meadow (SBO1), thicket-like (SBS1), or more closed and treed (SBT1). Tree cover always ≤ 60%. | Sand Barrens typically are exposed sand, generally sparsely vegetated and caused by lack of moisture, periodic fires and erosion. They have little or no soil and the underlying rock protrudes through the surface. Usually located within other types of natural habitat such as forest or savannah. Vegetation can vary from patchy and barren to tree covered but less than 60%. | A sand barren area >0.5ha in size <u>Information Sources</u> <ul style="list-style-type: none">• OMNRF Districts• Natural Heritage Information Centre (NHIC) has location information available on their website• Field naturalist clubs• Conservation Authorities | <ul style="list-style-type: none">• Confirm any ELC Vegetation Type for Sand Barrens^{lxxviii}• Site must not be dominated by exotic or introduced species (<50% vegetative cover are exotics sp)ⁱ.• SWHMIST^{cxlix} Index #20 provides development effects and mitigation measures. | Sand Barrens are not present within or adjacent to the subject property. | |

Table 2. Characteristics of Rare Vegetation Communities for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|--|--|--|--|--|------------------|
| | ELC Ecosite Codes | Habitat Description | Detailed Information and Sources | Defining Criteria | Study Area | Subject Property |
| Alvar | | | | | Not Present | Not Present |
| Alvars are extremely rare habitats in Ecoregion 7E | ALO1 ALS1 ALT1 FOC1 FOC2 CUM2 CUS2 CUT2-1 CUW2 Five Alvar Indicator Species: 1) <i>Carex crawei</i> 2) <i>Panicum philadelphicum</i> 3) <i>Eleocharis compressa</i> 4) <i>Scutellaria parvula</i> 5) <i>Trichostema brachiatum</i> These indicator species are very specific to Alvars within Ecoregion 7E ^{cxlix} | An alvar is typically a level, mostly unfractured calcareous bedrock feature with a mosaic of rock pavements and bedrock overlain by a thin veneer of soil. The hydrology of alvars is complex, with alternating periods of inundation and drought. Vegetation cover varies from sparse lichen-moss associations to grasslands and shrublands and comprising a number of characteristic or indicator plant. Undisturbed alvars can be phyto- and zoogeographically diverse, supporting many uncommon or are relict plant and animals species. Vegetation cover varies from patchy to barren with a less than 60% tree cover ^{lxxviii} . | An Alvar site > 0.5ha in size ^{lxxv} . Alvar is particularly rare in Ecoregion 7E where the only known sites are found in the western islands of Lake Erie ^{cxci} . <u>Information Sources</u> <ul style="list-style-type: none">Alvars of Ontario (2000), Federation of Ontario Naturalists^{lxxvi}.Ontario Nature – Conserving Great Lakes Alvars^{ccviii}.Natural Heritage Information Centre (NHIC) has location information available on their websiteOMNRF StaffField Naturalist clubsConservation Authorities | Field studies identify four of the five Alvar indicator species ^{lxxv} at a candidate Alvar site is Significant <ul style="list-style-type: none">Site must not be dominated by exotic or introduced species (<50% vegetative cover exotics).The alvar must be in excellent condition and fit in with surrounding landscape with few conflicting land uses^{lxxv}.SWHMIST^{cxlix} Index #17 provides development effects and mitigation measures. | Alvars are not present within or adjacent to the subject property. | |
| Old Growth Forest | | | | | Not Present | Not Present |
| Due to historic logging practices and land clearance for agriculture, old growth forest is rare in Ecoregion 7E. | Forest Community Series: FOD FOC FOM SWD SWC SWM | Old growth forests are characterized by heavy mortality or turnover of overstorey trees resulting in a mosaic of gaps that encourage development of a multi-layered canopy and an abundance of snags and downed woody debris. | Woodland area is >0.5ha <u>Information Sources</u> <ul style="list-style-type: none">OMNRF Forest Resource Inventory mappingOMNRF DistrictsField naturalist clubsConservation AuthoritiesSustainable Forestry Licence (SFL) companies will possibly know locations through field operations.Municipal forestry departments | Field Studies will determine: <ul style="list-style-type: none">If dominant trees species of the ecosite are >140 years old, then stand is Significant Wildlife Habitat^{cxlviii}.The forested area containing the old growth characteristics will have experienced no recognizable forestry activities^{cxlviii} (cut stumps will not be present)Determine ELC Vegetation Type for forest area containing the old growth characteristics^{lxxviii}.SWHMIST^{cxlix} Index #23 provides development effects and mitigation measures | Old growth forests are not present within or adjacent to the subject property. | |

Table 2. Characteristics of Rare Vegetation Communities for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|--|--|--|---|--|------------------|
| | ELC Ecosite Codes | Habitat Description | Detailed Information and Sources | Defining Criteria | Study Area | Subject Property |
| Savannah | | | | | Not Present | Not Present |
| Savannahs are extremely rare habitats in Ontario. | TPS1 TPS2 TPW1 TPW2 CUS2 | <p>A Savannah is a tallgrass prairie habitat that has tree cover between 25 – 60%.</p> <p>In Ecoregion 7E, known Tallgrass Prairie and savannah remnants are scattered between Lake Huron and Lake Erie, near Lake St. Clair, north of and along the Lake Erie shoreline, in Brantford and in the Toronto area (north of Lake Ontario)^{cc}.</p> | <p>No minimum size to site^l. Site must be restored or a natural site. Remnant sites such as railway right of ways are not considered to be SWH.</p> <p><u>Information Sources</u></p> <ul style="list-style-type: none">• OMNRF Districts• Natural Heritage Information Centre (NHIC) has location data available on their website• Field naturalists clubs• Conservation Authorities | <p>Field studies confirm one or more of the Savannah indicator species listed in^{lxxv} Appendix N should be present^l. Note: Savannah plant spp. list from Ecoregion 7E should be used.</p> <ul style="list-style-type: none">• Area of the ELC Vegetation type is the SWH^{lxxviii}.• Site must not be dominated by exotic or introduced species (<50% vegetative cover exotics).• SWHMIST^{cxlix} Index #18 provides development effects and mitigation measures. | Savannahs (i.e., tallgrass prairie habitat that has 25-60% tree cover) are not present within or adjacent to the subject property. | |
| Tallgrass Prairie | | | | | Not Present | Not Present |
| Tallgrass Prairies are extremely rare habitats in Ontario. | TPO1 TPO2 | <p>A Tallgrass Prairie has ground cover dominated by prairie grasses. An open Tallgrass Prairie habitat has < 25% tree cover.</p> <p>In Ecoregion 7E, known Tallgrass Prairie and savannah remnants are scattered between Lake Huron and Lake Erie, near Lake St. Clair, north of and along the Lake Erie shoreline, in Brantford and in the Toronto area (north of Lake Ontario)^{cc}.</p> | <p>No minimum size to site^l. Site must be restored or a natural site. Remnant sites such as railway right of ways are not considered to be SWH.</p> <p><u>Information Sources</u></p> <ul style="list-style-type: none">• Natural Heritage Information Centre (NHIC has location information available on their website• OMNRF Districts• Field naturalists clubs• Conservation Authorities | <p>Field studies confirm one or more of the Prairie indicator species listed in^{lxxv} Appendix N should be present^l. Note: Prairie plant spp. list from Ecoregion 7E should be used.</p> <ul style="list-style-type: none">• Area of the ELC Vegetation Type is the SWH^{lxxviii}.• Site must not be dominated by exotic or introduced species (<50% vegetative cover exotics).• SWHMIST^{cxlix} Index #19 provides development effects and mitigation measures. | Tallgrass Prairie habitat is not present within or adjacent to the subject property. | |
| Other Rare Vegetation Communities | | | | | Not Present | Not Present |
| Plant communities that often contain rare species which depend on the habitat for survival. | Provincially Rare S1, S2 and S3 vegetation communities are listed in Appendix M of the SWHTG ^{cxlviii} . Any ELC Ecosite Code that has a possible ELC Vegetation Type that is Provincially Rare is Candidate SWH. | Rare Vegetation Communities may include beaches, fens, forest, marsh, barrens, dunes and swamps. | <p>ELC Ecosite codes that have the potential to be a rare ELC Vegetation Type as outlined in appendix M^{cxlviii}.</p> <p>The OMNRF/NHIC will have up to date listing for rare vegetation communities.</p> <p><u>Information Sources</u></p> <ul style="list-style-type: none">• Natural Heritage Information Centre (NHIC) has location information available on their website• OMNRF Districts• Field naturalists clubs• Conservation Authorities | <p>Field studies should confirm if an ELC Vegetation Type is a rare vegetation community based on listing within Appendix M of SWHTG^{cxlviii}.</p> <ul style="list-style-type: none">• Area of the ELC Vegetation Type polygon is the SWH.• SWHMIST^{cxlix} Index #37 provides development effects and mitigation measures. | Beaches, fens, barrens, dunes, and swamps are not present within the subject property. Rare vegetation communities are not present within or adjacent to the subject property. | |

Significant Wildlife Habitat Assessment Tables

Table 3. Characteristics of Specialized Wildlife Habitat for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|--|--|--|--|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Waterfowl Nesting Area | | | | | Not Present | Not Present |
| Important to local waterfowl populations, sites with greatest number of species and highest number of individuals are significant | American Black Duck Northern Pintail Northern Shoveler Gadwall Blue-winged Teal Green-winged Teal Wood Duck Hooded Merganser Mallard | All upland habitats located adjacent to these wetland ELC Ecosites are Candidate SWH: <div>MAS1 MAS2 MAS3 SAS1 SAM1 SAF1 MAM1 MAM2 MAM3 MAM4 MAM5 MAM6 SWT1 SWT2 SWD1 SWD2 SWD3 SWD4</div> Note: includes adjacency to Provincially Significant Wetlands | A waterfowl nesting area extends: 120m ^{cxlix} from a wetland (>0.5ha) or a wetland (>0.5ha) with small wetlands (0.5ha) within 120m or a cluster of 3 or more small (<0.5 ha) wetlands within 120m of each individual wetland where waterfowl nesting is known to occur ^{cxlix} . • Upland areas should be at least 120m wide so that predators such as racoons, skunks, and foxes have difficulty finding nests. • Wood Ducks and Hooded Mergansers utilize large diameter trees (>40cm dbh) in woodlands for cavity nest sites. <u>Information Sources</u> • Ducks Unlimited staff may know the locations of particularly productive nesting sites. • OMNRF Wetland Evaluations for indication of significant waterfowl nesting habitat. • Reports and other information available from CAs | Studies confirmed: • Presence of 3 or more nesting pairs for listed species excluding Mallards ⁱ , or, • Presence of 10 or more nesting pairs for listed species including Mallards ⁱ . • Any active nesting site of an American Black Duck is considered significant. • Nesting studies should be completed during the spring breeding season (April - June). Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} • A field study confirming waterfowl nesting habitat will determine the boundary of the waterfowl nesting habitat for the SWH, this may be greater or less than 120m ^{cxlviii} from the wetland and will provide enough habitat for waterfowl to successfully nest. • SWHMIST ^{cxlix} Index #25 provides development effects and mitigation measures. | No wetlands within 120m are present within or adjacent to the subject property. | |
| Wildlife Habitat: Bald Eagle and Osprey Nesting, Foraging and Perching Habitat | | | | | Not Present | Not Present |
| Nest sites are fairly uncommon in Ecoregion 7E and are used annually by these species. Many suitable nesting locations may be lost due to increasing shoreline development pressures and scarcity of habitat. | Osprey <u>Special Concern:</u> Bald Eagle | ELC Forest Community Series: FOD, FOM, FOC, SWD, SWM and SWC directly adjacent to riparian areas – rivers, lakes, ponds and wetlands. | Nests are associated with lakes, ponds, rivers or wetlands along forested shorelines, islands, or on structures over water. Osprey nests are usually at the top a tree whereas Bald Eagle nests are typically in super canopy trees in a notch within the tree’s canopy. Nests located on man-made objects are not to be included as SWH (e.g. telephone poles and constructed nesting platforms). <u>Information Sources</u> • Natural Heritage Information Center (NHIC) compiles all known nesting sites for Bald Eagles in Ontario • MNRF values information (LIO/NRVIS) will list known nesting locations, Note: data from NRVIS is provided as a point format and does not include all the habitat. • Nature Counts, Ontario Nest Records Scheme data • OMNRF Districts • Check the Ontario Breeding Bird Atlas ^{ccv} or Rare Breeding Birds in Ontario for species documented • Reports and other information available from CAs • Field naturalists clubs | Studies confirm the use of these nests by: • One or more active Osprey or Bald Eagle nests in an area ^{cxlviii} . • Some species have more than one nest in a given area and priority is given to the primary nest with alternate nests included within the area of the SWH. • For an Osprey, the active nest and a 300m radius around the nest or the contiguous woodland stand is the SWH ^{ccvii} , maintaining undisturbed shorelines with large trees within this area is important ^{cxlviii} . • For a Bald Eagle the active nest and a 400-800m radius around the nest is the SWH ^{cvi, ccvii} . Area of the habitat from 400-800m is dependant on site lines from the nest to the development and inclusion of perching and foraging habitat ^{cvi} . • To be significant a site must be used annually. When found inactive, the site must be known to be inactive for ≥3 years or suspected of not being used for >5 years before being considered not significant ^{ccvii} . • Observational studies to determine nest site use, perching sites and foraging areas need to be done from mid March to mid August. • Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} • SWHMIST ^{cxlix} Index #26 provides development effects and mitigation measures. | Lakes, ponds, rivers, and wetlands are not present. Suitable habitat is not present within or adjacent to the subject property. | |

Table 3. Characteristics of Specialized Wildlife Habitat for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|---|--|---|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Woodland Raptor Nesting Habitat | | | | | Not Present | Not Present |
| Nests sites for these species are rarely identified; these area sensitive habitats are often used annually by these species. | Northern Goshawk Cooper’s Hawk Sharp-shinned Hawk Red-shouldered Hawk Barred Owl Broad-winged Hawk | May be found in all forested ELC Ecosites. May also be found in SWC, SWM, SWD and CUP3 | All natural or conifer plantation woodland/forest stands combined >30ha or with >4ha of interior habitat ^{lxxxviii, lxxxix, xc, xci, xciii, xciv, xcv,xcvi, cxxxiii} . Interior habitat determined with a 200m buffer ^{cxlviii} . • Stick nests found in a variety of intermediate-aged to mature conifer, deciduous or mixed forests within tops or crotches of trees. Species such as Coopers hawk nest along forest edges sometimes on peninsulas or small off-shore islands. • In disturbed sites, nests may be used again, or a new nest will be in close proximity to old nest. <u>Information Sources</u> • OMNRF Districts • Check the Ontario Breeding Bird Atlas ^{ccv} or Rare Breeding Birds in Ontario for species documented. • Check data from Bird Studies Canada • Reports and other information available from CAs | Studies confirm: • Presence of 1 or more active nests from species list is considered significant ^{cxlviii} . • Red-shouldered Hawk and Northern Goshawk – A 400m radius around the nest or 28 ha of habitat is the SWH ^{ccvii} . (the 28ha habitat area would be applied where optimal habitat is irregularly shaped around the nest) • Barred Owl – A 200m radius around the nest is the SWH ^{ccvii} . • Broad-winged Hawk and Coopers Hawk – A 100m radius around the nest is the SWH ^{ccvii} . • Sharp-Shinned Hawk – A 50m radius around the nest is the SWH ^{ccvii} . • Conduct field investigations from early March to end of May. The use of call broadcasts can help in locating territorial (courting/nesting) raptors and facilitate the discovery of nests by narrowing down the search area. • SWHMIST ^{cxlix} Index #27 provides development effects and mitigation measures. | The woodland does not meet the >30ha size criterion for this SWH. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Turtle Nesting Area | | | | | Not Present | Not Present |
| These habitats are rare and when identified will often be the only breeding site for local populations of turtles. | Midland Painted Turtle <u>Special Concern:</u> Northern Map Turtle Snapping Turtle | Exposed mineral soil (sand or gravel) areas adjacent (<100m) ^{cxlviii} or within the following ELC Ecosites: MAS1 MAS2 MAS3 SAS1 SAM1 SAF1 BOO1 FEO1 | • Best nesting habitat for turtles are close to water and away from roads and sites less prone to loss of eggs by predation from skunks, raccoons or other animals. • For an area to function as a turtle-nesting area, it must provide sand and gravel that turtles are able to dig in and are located in open, sunny areas. Nesting areas on the sides of municipal or provincial road embankments and shoulders are not SWH. • Sand and gravel beaches adjacent to undisturbed shallow weedy areas of marshes, lakes, and rivers are most frequently used. <u>Information Sources</u> • Use Ontario Soil Survey reports and maps to help find suitable substrate for nesting turtles (well-drained sands and fine gravels). • Check the Ontario Herpetofaunal Summary Atlas records or other similar atlases for uncommon turtles; location information may help to find potential nesting habitat for them. • Natural Heritage Information Center (NHIC) Field naturalist clubs | Studies confirm: • Presence of 5 or more nesting Midland Painted Turtles ^l • One or more Northern Map Turtle or Snapping Turtle nesting is a SWH ^l • The area or collection of sites within an area of exposed mineral soils where the turtles nest, plus a radius of 30-100m around the nesting area dependant on slope, riparian vegetation and adjacent land use is the SWH ^{cxlviii} . • Travel routes from wetland to nesting area are to be considered within the SWH as part of the 30-100m area of habitat ^{cxlix} . • Field investigations should be conducted in prime nesting season typically late spring to early summer. Observation studies observing the turtles nesting is a recommended method. • SWHMIST ^{cxlix} Index #28 provides development effects and mitigation measures for turtle nesting habitat. | Adjacent areas are highly disturbed (i.e., recreational fields with human presence, surrounding roads), do not contain sandy areas and are not near water. Suitable habitat is not present within or adjacent to the subject property. | |

Table 3. Characteristics of Specialized Wildlife Habitat for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|---|---|--|--|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Seeps and Springs | | | | | Not Present | Not Present |
| Seeps/Springs are typical of headwater areas and are often at the source of coldwater streams | Wild Turkey Ruffed Grouse Spruce Grouse White-tailed Deer Salamander spp. | Seeps/Springs are areas where ground water comes to the surface. Often they are found within headwater areas within forested habitats. Any forested Ecosite within the headwater areas of a stream could have seeps/springs. | Any forested area (with <25% meadow/field/pasture) within the headwaters of a stream or river system ^{cxvii, cxlix} . • Seeps and springs are important feeding and drinking areas especially in the winter will typically support a variety of plant and animal species ^{cxix, cxx, cxxi, cxxii, cxiii, cxiv} . <u>Information Sources</u> • Topographical Map • Thermography • Hydrological surveys conducted by CAs and MOE • Field naturalists and landowners • Municipalities and Conservation Authorities may have drainage maps and headwater areas mapped | Field Studies confirm: • Presence of a site with 2 or more ⁱ seeps/springs should be considered SWH. • The area of a ELC forest ecosite containing the seeps/springs is the SWH. The protection of the recharge area considering the slope, vegetation, height of trees and groundwater condition need to be considered in delineation of the habitat ^{cxlviii} . • SWHMIST ^{cxlix} Index #30 provides development effects and mitigation measures. | Seeps and springs are not present within or adjacent the subject property. The on-site Headwater Drainage Feature is largely formed by drainage from Munn's School. | |
| Wildlife Habitat: Amphibian Breeding Habitat (Woodland) | | | | | Not Present | Not Present |
| These habitats are extremely important to amphibian biodiversity within a landscape and often represent the only breeding habitat for local amphibian populations | Eastern Newt Blue-spotted Salamander Spotted Salamander Gray Treefrog Spring Peeper Western Chorus Frog Wood Frog | All Ecosites associated with these ELC Community Series: FOC FOM FOD SWC SWM SWD Breeding pools within the woodland or the shortest distance from forest habitat are more significant because they are more likely to be used due to reduced risk to migrating amphibians. | • Presence of a wetland, pond or woodland pool (including vernal pools) >500m ² (about 25m diameter) ^{ccvii} within or adjacent (within 120m) to a woodland (no minimum size) ^{clxxxii, lxiii, lxv, lxvi, lxvii, lxviii, lxix, lxx} . Some small wetlands may not be mapped and may be important breeding pools for amphibians. • Woodlands with permanent ponds or those containing water in most years until mid-July are more likely to be used as breeding habitat ^{cxlviii} . <u>Information Sources</u> • Ontario Herpetofaunal Summary Atlas (or other similar atlases) for records • Local landowners may also provide assistance as they may hear spring-time choruses of amphibians on their property. • OMNRF Districts and wetland evaluations • Field naturalist clubs • Canadian Wildlife Service Amphibian Road Call Survey • Ontario Vernal Pool Association: http://www.ontariovernalpools.org | Studies confirm: • Presence of breeding population of 1 or more of the listed newt/salamander species or 2 or more of the listed frog/toad species with at least 20 individuals (adults or eggs masses) or 2 or more of the listed frog/toad species with Call Level Codes of 3. • A combination of observational study and call count surveys ^{cviii} will be required during the spring (March-June) when amphibians are concentrated around suitable breeding habitat within or near the woodland/wetlands. • The habitat is the wetland area plus a 230m radius of woodland area ^{lxiii, lxv, lxvi, lxvii, lxviii, lxix, lxx, lxxi} . If a wetland area is adjacent to a woodland, a travel corridor connecting the wetland to the woodland is to be included in the habitat. • SWHMIST ^{cxlix} Index #14 provides development effects and mitigation measures. | Wetlands, ponds, and woodland pools (including vernal pools) are not located adjacent (within 120m) to a woodland. Suitable habitat is not present within or adjacent to the subject propery | |

Table 3. Characteristics of Specialized Wildlife Habitat for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|---|---|--|---|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Amphibian Breeding Habitat (Wetland) | | | | | Not Present | Not Present |
| Wetlands supporting breeding for these amphibian species are extremely important and fairly rare within Central Ontario Landscapes | Eastern Newt American Toad Spotted Salamander Four-toed Salamander Blue-spotted Salamander Gray Treefrog Western Chorus Frog Northern Leopard Frog Pickerel Frog Green Frog Mink Frog Bullfrog | ELC Community Classes SW, MA, FE, BO, OA and SA. Typically these wetland ecosites will be isolated (>120m) from woodland ecosites, however larger wetlands containing predominantly aquatic species (e.g. Bull Frog) may be adjacent to woodlands. | <ul style="list-style-type: none">Wetlands >500m² (about 25m diameter)^{ccvii} supporting high species diversity are significant: some small or ephemeral habitats may not be identified on MNR mapping and could be important amphibian breeding habitats^{clxxxiv}.Presence of shrubs and logs increase significance of pond for some amphibian species because of available structure for calling, foraging, escape and concealment from predators.Bullfrogs require permanent water bodies with abundant emergent vegetation. <u>Information Sources</u> <ul style="list-style-type: none">Ontario Herpetofaunal Summary Atlas (or other similar atlases)Canadian Wildlife Service Amphibian Road Surveys and Backyard Amphibian Call Count.OMNRF Districts and wetland evaluationsReports and other information available from CAs | Studies confirm: <ul style="list-style-type: none">Presence of breeding population of 1or more of the listed newt/salamander species or 2 or more of the listed frog or toad species and with at least 20 breeding individuals (adults and eggs masses)^{lxxi, lxxiii} or 2 or more of the listed frog/toad species with Call Level of 3. or; Wetland with confirmed breeding Bullfrogs are significant^l.The ELC ecosite wetland area and the shoreline are the SWH.A combination of observational study and call count surveys cviii to determine breeding/larval stages will be required during the spring (May March-June) when amphibians are concentrated around suitable breeding habitat within or near the woodland/wetlands.If a SWH is determined for Amphibian Breeding Habitat (Wetlands) then Movement Corridors are to be considered as outlined in Table 1.4.1 of this Schedule.SWHMIST^{cdlix} Index #15 provides development effects and mitigation measures. | Wetlands are not present within or adjacent to the subject property. | |
| Wildlife Habitat: Woodland Area-Sensitive Bird Breeding Habitat | | | | | Not Present | Not Present |
| Large, natural blocks of mature woodland habitat within the settled areas of Southern Ontario are important habitats for area sensitive interior forest song birds. | Yellow-bellied Sapsucker Red-breasted Nuthatch Veery Blue-headed Vireo Northern Parula Black-throated Green Warbler Blackburnian Warbler Black-throated Blue Warbler Ovenbird Scarlet Tanager Winter Wren Pileated Woodpecker <u>Special Concern:</u> Cerulean Warbler Canada Warbler | All Ecosites associated with these ELC Community Series: FOC FOM FOD SWC SWM SWD | <ul style="list-style-type: none">Habitats where interior forest breeding birds are breeding, typically large mature (>60 yrs. old) forest stands or woodlots >30ha^{cv, cxxxi, cxxxii, cxxxiii, cxxxiv, cxxxv, cxxxvi, cxxxvii, cxxxviii, cxxxix, cxl, cxli, cxlii, cxliii, cxliv, cxlv, cxlvi, cli, clii, cliii, cliv, clv, clvi, clvii, clviii, clix}.Interior forest habitat is at least 200m from forest edge habitat^{clxiv}. <u>Information Sources</u> <ul style="list-style-type: none">Local birder clubsCanadian Wildlife Service (CWS) for the location of forest bird monitoringBird Studies Canada conducted a 3-year study of 287 woodlands to determine the effects of forest fragmentation on forest birds and to determine what forests were of greatest value to interior species.Reports and other information available from CAs | Studies confirm: <ul style="list-style-type: none">Presence of nesting or breeding pairs of 3 or more of the listed wildlife species^l.Note: any site with breeding Cerulean Warblers or Canada Warbler is to be considered SWH^l.Conduct field investigations in early summer when birds are singing and defending their territories.Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects”^{ccxi}SWHMIST^{cdlix} Index #34 provides development effects and mitigation measures. | The on-site woodland does not meet the >30ha size criterion for this SWH. Further, large mature (>60yrs old) forest stands or woodlots are not present. Suitable habitat is not present within or adjacent to the subject property. | |

Significant Wildlife Habitat Assessment Tables

Table 4. Characteristics of Habitat for Species of Conservation Concern for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|---|--|---|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Marsh Bird Breeding Habitat | | | | | Not Present | Not Present |
| Wetlands for these bird species are typically productive and fairly rare in Southern Ontario landscapes. | American Bittern Virginia Rail Sora Common Gallinule American Coot Pied-billed Grebe Marsh Wren Sedge Wren Common Loon Green Heron Trumpeter Swan <u>Special Concern:</u> Black Tern Yellow Rail | MAM1 MAM2 MAM3 MAM4 MAM5 MAM6 SAS1 SAM1 SAF1 FEO1 BOO1 For Green Heron: All SW, MA and CUM1 sites | <ul style="list-style-type: none">Nesting occurs in wetlandsAll wetland habitat is to be considered as long as there is shallow water with emergent aquatic vegetation present^{cxxiv}.For Green Heron, habitat is at the edge of water such as sluggish streams, ponds and marshes sheltered by shrubs and trees. Less frequently, it may be found in upland shrubs or forest a considerable distance from water. <u>Information Sources</u> <ul style="list-style-type: none">OMNRF Districts and wetland evaluationsField naturalist clubsNatural Heritage Information Centre (NHIC)Reports and other information available from CAsOntario Breeding Bird Atlas^{ccv} | Studies confirm: <ul style="list-style-type: none">Presence of 5 or more nesting pairs of Sedge Wren or Marsh Wren or breeding by any combination of 4 or more of the listed species^l.Note: any wetland with breeding of 1 or more Trumpeter Swans, Black Terns, Green Heron or Yellow Rail is SWH^l.Area of the ELC ecosite is the SWHBreeding surveys should be done in May/June when these species are actively nesting in wetland habitats.Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects”^{ccxi}SWHMIST^{cxlix} Index #35 provides development effects and mitigation measures | Wetlands are not present within or adjacent to the subject property. | |
| Wildlife Habitat: Open Country Bird Breeding Habitat | | | | | Not Present | Not Present |
| This wildlife habitat is declining throughout Ontario and North America. Species such as the Upland Sandpiper have declined significantly the past 40 years based on CWS (2004) trend records. | Upland Sandpiper Grasshopper Sparrow Vesper Sparrow Northern Harrier Savannah Sparrow <u>Special Concern:</u> Short-eared Owl | CUM1 CUM2 | <p>Large grassland areas (includes natural and cultural fields and meadows) >30ha^{clx, clxi, clxii, clxiii, clxiv, clxv, clxvi, clxvii, clxviii, clxix}.</p> <p>Grasslands not Class 1 or 2 agricultural lands, and not being actively used for farming (i.e. no row cropping or intensive hay or livestock pasturing in the last 5 years)^l.</p> <p>Grassland sites considered significant should have a history of longevity, either abandoned fields, mature hayfields and pasturelands that are at least 5 years or older.</p> <p>The Indicator bird species are area sensitive requiring larger grassland areas than the common grassland species.</p> <u>Information Sources</u> <ul style="list-style-type: none">Agricultural land classification maps Ministry of AgricultureLocal birder clubsOntario Breeding Bird Atlas^{ccv}EIS Reports and other information available from CAs | Field Studies confirm: <ul style="list-style-type: none">Presence of nesting or breeding of 2 or more of the listed species^l.A field with 1 or more breeding Short-eared Owls is to be considered SWH.The area of SWH is the contiguous ELC ecosite field areas.Conduct field investigations of the most likely areas in spring and early summer when birds are singing and defending their territories.Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects”^{ccxi}SWHMIST^{cxlix} Index #32 provides development effects and mitigation measures | Large grassland areas (>30ha in size) are not present within or adjacent to the subject property. | |

Table 4. Characteristics of Habitat for Species of Conservation Concern for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|---|--|--|---|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Shrub/Early Successional Bird Breeding Habitat | | | | | Not Present | Not Present |
| This wildlife habitat is declining throughout Ontario and North America. The Brown Thrasher has declined significantly over the past 40 years based on CWS (2004) trend records. | Indicator Spp: Brown Thrasher Clay-coloured Sparrow Common Spp. Field Sparrow Black-billed Cuckoo Eastern Towhee Willow Flycatcher <u>Special Concern:</u> Yellow-breasted Chat Golden-winged Warbler | CUT1 CUT2 CUS1 CUS2 CUW1 CUW2 Patches of shrub ecosites can be complexed into a larger habitat such as woodland area for some bird species. | Large natural field areas succeeding to shrub and thicket habitats >10ha ^{clxiv} in size. Shrub land or early successional fields, not class 1 or 2 agricultural lands, not being actively used for farming (i.e. no row-cropping, haying or live-stock pasturing in the last 5 years) ^l . Shrub thicket habitats (>10 ha) are most likely to support and sustain a diversity of these species ^{clxxiii} . Shrub and thicket habitat sites considered significant should have a history of longevity, either abandoned fields or pasturelands. <u>Information Sources</u> • Agricultural land classification maps, Ministry of Agriculture. • Local bird clubs • Ontario Breeding Bird Atlas ^{ccv} • Reports and other information available from CAs | Field Studies confirm: • Presence of nesting or breeding of 1 of the indicator species and at least 2 of the common species ^l . • A field with breeding Yellow-breasted Chat or Golden-winged Warbler is to be considered as Significant Wildlife Habitat ^l . • The area of the SWH is the contiguous ELC ecosite field/thicket area. • Conduct field investigations of the most likely areas in spring and early summer when birds are singing and defending their territories • Evaluation methods to follow “Bird and Bird Habitats: Guidelines for Wind Power Projects” ^{ccxi} • SWHMIST ^{cxlix} Index #33 provides development effects and mitigation measures. | Large natural field areas succeeding to shrub and thicket habitats >10ha in size are not present. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Terrestrial Crayfish | | | | | Not Present | Not Present |
| Terrestrial Crayfish are only found within SW Ontario in Canada and their habitats are very rare. ^{Ccii} | Chimney or Digger Crayfish (<i>Fallicambarus fodiens</i>) Devil Crawfish or Meadow Crayfish (<i>Cambarus Diogenes</i>) | MAM1 MAM2 MAM3 MAM4 MAM5 MAM6 MAS1 MAS2 MAS3 SWD SWT SWM CUM1 with inclusions of above meadow marsh ecosites can be used by terrestrial crayfish. | Wet meadow and edges of shallow marshes (no minimum size) identified should be surveyed for terrestrial crayfish. • Constructs burrows in marshes, mudflats, meadows, the ground can’t be too moist. Can often be found far from water. • Both species are a semi-terrestrial burrower which spends most of its life within burrows consisting of a network of tunnels. Usually the soil is not too moist so that the tunnel is well formed. <u>Information Sources</u> • Information sources from “Conservation Status of Freshwater Crayfishes” by Dr. Premek Hamr for the WWF and CNF March 1998. | Studies Confirm: • Presence of 1 or more individuals of species listed or their chimneys (burrows) in suitable marsh meadow or terrestrial sites ^{cci} . • Area of ELC Ecosite or an ecoelement area of meadow marsh or swamp within the large ecosite area is the SWH • Surveys should be done April to August in temporary or permanent water. Note the presence of burrows or chimneys are often the only indicator of presence, observance or collection of individuals is very difficult ^{cci} • SWHMIST ^{cxlix} Index #36 provides development effects and mitigation measures. | Wet meadows and edges of shallow marshes are not present. Suitable habitat is not present within or adjacent to the subject property. | |
| Wildlife Habitat: Special Concern and Rare Wildlife Species | | | | | Not Present | Not Present |
| These species are quite rare or have experienced significant population declines in Ontario | All Special Concern and Provincially Rare (S1-S3, SH) plant and animal species. Lists of these species are tracked by the Natural Heritage Information Centre (NHIC). | All plant and animal element occurrences (EO) within a 1 or 10km grid. Older element occurrences were recorded prior to GPS being available, therefore location information may lack accuracy. | When an element occurrence is identified within a 1 or 10 km grid for a Special Concern or provincially Rare species; linking candidate habitat on the site needs to be completed to ELC Ecosites ^{lxxviii} . <u>Information Sources</u> • Natural Heritage Information Centre (NHIC) will have the Special Concern and Provincially Rare (S1-S3, SH) species lists and element occurrences for these species. • NHIC Website: "Get Information" http://nhic.mnr.gov.on.ca • Ontario Breeding Bird Atlas ^{ccv} • Expert advice should be sought as many of the rare spp. have little information available about their requirements. | Studies Confirm: • Assessment/inventory of the site for the identified special concern or rare species needs to be completed during the time of year when the species is present or easily identifiable. • The area of the habitat to the finest ELC scale that protects the habitat form and function is the SWH, this must be delineated through detailed field studies. The habitat neess to be easily mapped and cover an important life stage component for a species e.g. specific nesting habitat for foraging habitat. • SWHMIST ^{cxlix} Index #37 provides development effects and mitigation measures. | Suitable habitat for SCC is not present within or adjacent to the subject property. | |

Significant Wildlife Habitat Assessment Tables

Table 5. Characteristics of Animal Movement Corridors for Ecoregion 7E (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|---|--|---|--|---|--|------------------|
| | Wildlife Species | ELC Ecosite Codes | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Wildlife Habitat: Amphibian Movement Corridors | | | | | Not Present | Not Present |
| Movement corridors for amphibians moving from their terrestrial habitat to breeding habitat can be extremely important for local populations. | Eastern Newt American Toad Blue-spotted Salamander Spotted Salamander Four-toed Salamander Gray Treefrog Northern Leopard Frog Pickerel Frog Western Chorus Frog | Corridors may be found in all ecosites associated with water. • Corridors will be determined based on identifying the significant breeding habitat for these species in Table 1.1. | Movement corridors between breeding habitat and summer habitat ^{clxxiv, clxxv, clxxvi, clxxvii, clxxviii, clxxix, clxxx, clxxxi} Movement corridors must be considered when Amphibian breeding habitat is confirmed as SWH from Table 1.2.2 (Amphibian Breeding Habitat – Wetland) of this Schedule ⁱ . <u>Information Sources</u> • MNRF District Office • Natural Heritage Information Centre NHIC • Reports and other information available from CAs • Field naturalist Clubs | • Field Studies must be conducted at the time of year when species are expected to be migrating or entering breeding sites. • Corridors should consist of native vegetation, with several layers of vegetation. Corridors unbroken by roads, waterways or bodies, and undeveloped areas are most significant ^{cxlix} . • Corridors should have at least 15m of vegetation on both sides of waterway ^{cxlix} or be up to 200m wide ^{cxlix} of woodland habitat and with gaps <20m ^{cxlix} • Shorter corridors are more significant than longer corridors, however amphibians must be able to get to and from their summer and breeding habitat ^{cxlix} . • SWHMIST ^{cxlix} Index #40 provides development effects and mitigation measures. | Amphibian breeding habitat is not present within and adjacent to the subject property (i.e., wetlands are not present). Therefore, suitable amphibian movement corridors are not present within or adjacent to the subject property. | |

Significant Wildlife Habitat Assessment Tables

Table 6. Exceptions for Ecodistricts within Ecoregion 7E-2 (MNRF 2015)

| Rationale | Candidate SWH | | | Confirmed SWH | Assessment Details | |
|--|---|-----------------------|---|--|---|------------------|
| | Wildlife Species | Ecosites | Habitat Criteria and Information Sources | Defining Criteria | Study Area | Subject Property |
| Bat Migratory Stopover Area | | | | | Not Present | Not Present |
| Stopover areas for long distance migrant bats are important during fall migration. | Hoary Bat Eastern Red Bat Silver-haired Bat | No specific ELC types | <ul style="list-style-type: none"> Long distance migratory bats typically migrate during late summer and early fall migrating summer breeding habitats throughout Ontario to southern wintering areas. Their annual fall migration may concentrate these species of bats at stopover areas. This is the only known bat migratory stopover habitats based on current information. <p><u>Information Sources</u></p> <ul style="list-style-type: none"> OMNRF for possible locations and contact for local experts University of Waterloo, Biology Department | <ul style="list-style-type: none"> Long Point (42°35'N, 80°30'E, to 42°33'N, 80°03'E) has been identified as a significant stop-over habitat for fall migrating Silver-haired bats, due to significant increases in abundance, activity and feeding that was documented during fall migration^{ccxv}. The confirmation criteria and habitat areas for this SWH are still being determined. SWHMIST^{cxlix} Index #38 provides development effects and mitigation measures | <p>The study area is not near Long Point. Stopover areas are not known to be present within the study area. Suitable habitat is not present within or adjacent to the subject property.</p> | |

Appendix III
Vascular Flora Species Observed within the Subject Property

Plant Species Reported from the Study Area - 1493 Sixth Line, Oakville (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Halton Region | NRSI Observed |
|-----------------------------------|----------------------------------|----------|-----------|---------------------------|---------------------------|---------------------------|---------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Varga 2000 | NRSI Results From 2023 and 2025 |
| Dicotyledons | Dicots | | | | | | | |
| Aceraceae | Maple Family | | | | | | | |
| <i>Acer negundo</i> | Manitoba Maple | S5 | | | | | X | X |
| <i>Acer platanoides</i> | Norway Maple | SE5 | | | | | X | X |
| <i>Acer saccharinum</i> | Silver Maple | S5 | | | | | X | X |
| Anacardiaceae | Sumac or Cashew Family | | | | | | | |
| <i>Rhus typhina</i> | Staghorn Sumac | S5 | | | | | X | X |
| Asteraceae | Composite or Aster Family | | | | | | | |
| <i>Ambrosia artemisiifolia</i> | Common Ragweed | S5 | | | | | X | X |
| <i>Arctium lappa</i> | Great Burdock | SE5 | | | | | X | X |
| <i>Arctium minus</i> | Common Burdock | SE5 | | | | | X | X |
| <i>Cirsium arvense</i> | Creeping Thistle | SE5 | | | | | X | X |
| <i>Cirsium vulgare</i> | Bull Thistle | SE5 | | | | | X | X |
| <i>Erigeron philadelphicus</i> | Philadelphia Fleabane | S5 | | | | | X | X |
| <i>Lapsana communis</i> | Common Nipplewort | SE5 | | | | | X | X |
| <i>Solidago sp.</i> | Goldenrod sp. | | | | | | | X |
| <i>Solidago altissima</i> | Tall Goldenrod | S5 | | | | | | X |
| <i>Sonchus arvensis</i> | Field Sow-thistle | SE5 | | | | | | X |
| <i>Symphyotrichum sp.</i> | Aster sp. | | | | | | | X |
| <i>Symphyotrichum lanceolatum</i> | Panicked Aster | S5 | | | | | X | X |
| <i>Taraxacum officinale</i> | Common Dandelion | SE5 | | | | | X | X |
| Balsaminaceae | Touch-me-not Family | | | | | | | |
| <i>Impatiens capensis</i> | Spotted Jewelweed | S5 | | | | | X | X |
| Betulaceae | Birch Family | | | | | | | |
| <i>Betula alleghaniensis</i> | Yellow Birch | S5 | | | | | X | X |
| Bignoniaceae | Bignonia Family | | | | | | | |
| <i>Catalpa speciosa</i> | Northern Catalpa | SE1 | | | | | | X |
| Boraginaceae | Borage Family | | | | | | | |
| <i>Hackelia virginiana</i> | Virginia Stickseed | S5 | | | | | U | X |
| Brassicaceae | Mustard Family | | | | | | | |
| <i>Alliaria petiolata</i> | Garlic Mustard | SE5 | | | | | X | X |
| <i>Barbarea vulgaris</i> | Bitter Wintercress | SE5 | | | | | X | X |
| Caprifoliaceae | Honeysuckle Family | | | | | | | |
| <i>Lonicera tatarica</i> | Tatarian Honeysuckle | SE5 | | | | | X | X |
| <i>Sambucus canadensis</i> | Common Elderberry | S5 | | | | | X | X |
| Caryophyllaceae | Pink Family | | | | | | | |
| <i>Dianthus armeria</i> | Deptford Pink | SE5 | | | | | X | X |
| Chenopodiaceae | Goosefoot Family | | | | | | | |
| <i>Chenopodium album</i> | White Goosefoot | SE5 | | | | | X | X |
| Clusiaceae | St. John's-wort Family | | | | | | | |
| <i>Hypericum perforatum</i> | Common St. John's-wort | SE5 | | | | | X | X |
| Cornaceae | Dogwood Family | | | | | | | |
| <i>Cornus racemosa</i> | Gray Dogwood | S5 | | | | | X | X |
| Fabaceae | Pea Family | | | | | | | |
| <i>Medicago lupulina</i> | Black Medic | SE5 | | | | | X | X |
| <i>Robinia pseudoacacia</i> | Black Locust | SE5 | | | | | X | X |
| <i>Trifolium aureum</i> | Yellow Clover | SE5 | | | | | X | X |
| <i>Trifolium repens</i> | White Clover | SE5 | | | | | X | X |
| <i>Vicia cracca</i> | Tufted Vetch | SE5 | | | | | X | X |
| Geraniaceae | Geranium Family | | | | | | | |
| <i>Geranium robertianum</i> | Herb-Robert | S5 | | | | | X | X |
| Grossulariaceae | Currant Family | | | | | | | |
| <i>Ribes rubrum</i> | Northern Red Currant | SE5 | | | | | X | X |
| Juglandaceae | Walnut Family | | | | | | | |
| <i>Carya cordiformis</i> | Bitternut Hickory | S5 | | | | | X | X |
| <i>Carya ovata</i> | Shagbark Hickory | S5 | | | | | | X |

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Halton Region | NRSI Observed |
|----------------------------------|-------------------------------------|----------|-----------|---------------------------|---------------------------|---------------------------|---------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Varga 2000 | NRSI Results From 2023 and 2025 |
| <i>Juglans nigra</i> | Black Walnut | S4? | | | | | X | X |
| Lamiaceae | Mint Family | | | | | | | |
| <i>Glechoma hederacea</i> | Ground Ivy | SE5 | | | | | X | X |
| <i>Leonurus cardiaca</i> | Common Motherwort | SE5 | | | | | | X |
| <i>Nepeta cataria</i> | Catnip | SE5 | | | | | X | X |
| <i>Prunella vulgaris</i> | Self-heal | S5 | | | | | | X |
| Malvaceae | Mallow Family | | | | | | | |
| <i>Abutilon theophrasti</i> | Velvetleaf | SE5 | | | | | X | X |
| Oleaceae | Olive Family | | | | | | | |
| <i>Fraxinus americana</i> | White Ash | S4 | | | | | X | X |
| <i>Ligustrum vulgare</i> | European Privet | SE5 | | | | | X | X |
| Onagraceae | Evening-primrose Family | | | | | | | |
| <i>Circaea canadensis</i> | Broad-leaved Enchanter's Nightshade | S5 | | | | | X | X |
| <i>Oenothera sp.</i> | Evening-primrose sp. | | | | | | | X |
| Oxalidaceae | Wood Sorrel Family | | | | | | | |
| <i>Oxalis stricta</i> | Upright Yellow Wood-sorrel | SE5 | | | | | X | X |
| Plantaginaceae | Plantain Family | | | | | | | |
| <i>Plantago major</i> | Common Plantain | SE5 | | | | | X | X |
| Primulaceae | Primrose Family | | | | | | | |
| <i>Lysimachia nummularia</i> | Creeping Jennie | SE5 | | | | | X | X |
| Ranunculaceae | Buttercup Family | | | | | | | |
| <i>Ranunculus caricetorum</i> | Northern Swamp Buttercup | S5 | | | | | X | X |
| <i>Ranunculus pennsylvanicus</i> | Pennsylvania Buttercup | S5 | | | | | U | X |
| Rhamnaceae | Buckthorn Family | | | | | | | |
| <i>Rhamnus cathartica</i> | Common Buckthorn | SE5 | | | | | X | X |
| Rosaceae | Rose Family | | | | | | | |
| <i>Agrimonia gryposepala</i> | Hooked Agrimony | S5 | | | | | X | X |
| <i>Crataegus sp.</i> | Hawthorn sp. | | | | | | | X |
| <i>Crataegus punctata</i> | Dotted Hawthorn | S5 | | | | | X | X |
| <i>Fragaria virginiana</i> | Wild Strawberry | S5 | | | | | | X |
| <i>Geum sp.</i> | Avens sp. | | | | | | | X |
| <i>Geum canadense</i> | White Avens | S5 | | | | | X | X |
| <i>Geum urbanum</i> | Wood Avens | SE3 | | | | | X | X |
| <i>Potentilla recta</i> | Sulphur Cinquefoil | SE5 | | | | | X | X |
| <i>Prunus avium</i> | Sweet Cherry | SE4 | | | | | X | X |
| <i>Prunus serotina</i> | Black Cherry | S5 | | | | | X | X |
| <i>Prunus virginiana</i> | Choke Cherry | S5 | | | | | X | X |
| <i>Rosa multiflora</i> | Multiflora Rose | SE5 | | | | | X | X |
| <i>Rubus idaeus</i> | Common Red Raspberry | S5 | | | | | | X |
| <i>Rubus occidentalis</i> | Black Raspberry | S5 | | | | | X | X |
| Rubiaceae | Madder Family | | | | | | | |
| <i>Galium aparine</i> | Cleavers | S5 | | | | | U | X |
| <i>Galium asprellum</i> | Rough Bedstraw | S5 | | | | | U | X |
| Scrophulariaceae | Figwort Family | | | | | | | |
| <i>Veronica serpyllifolia</i> | Thyme-leaved Speedwell | SU | | | | | X | X |
| Solanaceae | Nightshade Family | | | | | | | |
| <i>Solanum dulcamara</i> | Bittersweet Nightshade | SE5 | | | | | X | X |
| Ulmaceae | Elm Family | | | | | | | |
| <i>Ulmus americana</i> | American Elm | S5 | | | | | X | X |
| <i>Ulmus pumila</i> | Siberian Elm | SE3 | | | | | X | X |
| Verbenaceae | Vervain Family | | | | | | | |
| <i>Verbena urticifolia</i> | White Vervain | S5 | | | | | X | X |
| Violaceae | Violet Family | | | | | | | |
| <i>Viola nephrophylla</i> | Northern Bog Violet | S5 | | | | | | X |
| <i>Viola odorata</i> | English Violet | SE2 | | | | | X | X |
| Vitaceae | Grape Family | | | | | | | |
| <i>Parthenocissus vitacea</i> | Thicket Creeper | S5 | | | | | X | X |
| <i>Vitis riparia</i> | Riverbank Grape | S5 | | | | | X | X |

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Halton Region | NRSI Observed |
|----------------------------|-----------------------|----------|-----------|---------------------------|---------------------------|---------------------------|---------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Varga 2000 | NRSI Results From 2023 and 2025 |
| Monocotyledons | Monocots | | | | | | | |
| Cyperaceae | Sedge Family | | | | | | | |
| <i>Carex vulpinoidea</i> | Fox Sedge | S5 | | | | | X | X |
| <i>Scirpus atrocinctus</i> | Black-girdled Bulrush | S5 | | | | | | X |
| Juncaceae | Rush Family | | | | | | | |
| <i>Juncus effusus</i> | Soft Rush | S5 | | | | | | X |
| Poaceae | Grass Family | | | | | | | |
| <i>Glyceria striata</i> | Fowl Mannagrass | S5 | | | | | X | X |
| <i>Poa pratensis</i> | Kentucky Bluegrass | S5 | | | | | | X |
| Total | | | | | | | | 84 |

References

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>

Varga, S., editor. August 2000. Distribution and status of the vascular plants of the Greater Toronto Area. Ontario Ministry of Natural Resources, Aurora District. 103 pp.

Appendix IV
Bird Species Report from the Study Area

Bird Species Reported from the Study Area - 1493 Sixth Sixth Line, Oakville (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | OBBA* | NHIC Data** | NRSI Observed: Highest Level of Breeding Evidence |
|----------------------------------|--|----------|-----------|---------------------------|---------------------------|---------------------------|-----------------|-------------|--|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | BSC et al. 2006 | MNR 2024 | NRSI Results from 2023 and 2025 |
| Anatidae | Ducks, Geese & Swans | | | | | | | | |
| <i>Aix sponsa</i> | Wood Duck | S5B,S3N | | | | | CO | | |
| <i>Anas platyrhynchos</i> | Mallard | S5 | | | | | CO | | |
| <i>Anas rubripes</i> | American Black Duck | S4 | | | | | PR | | |
| <i>Branta canadensis</i> | Canada Goose | S5 | | | | | CO | | |
| <i>Cygnus olor</i> | Mute Swan | SNA | | | | | CO | | |
| Odontophoridae | New World Quails | | | | | | | | |
| <i>Colinus virginianus</i> | Northern Bobwhite | S1? | END | E | E | Schedule 1 | | X | |
| Columbidae | Pigeons & Doves | | | | | | | | |
| <i>Columba livia</i> | Rock Pigeon | SNA | | | | | CO | | |
| <i>Zenaidura macroura</i> | Mourning Dove | S5 | | | | | CO | | PO |
| Cuculiformes | Cuckoos & Anis | | | | | | | | |
| <i>Coccyzus americanus</i> | Yellow-billed Cuckoo | S4B | | | | | PO | | |
| <i>Coccyzus erythrophthalmus</i> | Black-billed Cuckoo | S4S5B | | | | | CO | | |
| Caprimulgidae | Goatsuckers | | | | | | | | |
| <i>Anthus vociferus</i> | Eastern Whip-poor-will | S4B | THR | SC | T | Schedule 1 | PR | | |
| <i>Chordeiles minor</i> | Common Nighthawk | S4B | SC | SC | SC | Schedule 1 | PO | | |
| Apodidae | Swifts | | | | | | | | |
| <i>Chaetura pelagica</i> | Chimney Swift | S3B | THR | T | T | Schedule 1 | CO | X | OB |
| Trochilidae | Hummingbirds | | | | | | | | |
| <i>Archilochus colubris</i> | Ruby-throated Hummingbird | S5B | | | | | PO | | |
| Charadriidae | Plovers & Lapwings | | | | | | | | |
| <i>Charadrius vociferus</i> | Killdeer | S4B | | | | | CO | | |
| Scolopacidae | Sandpipers & Allies | | | | | | | | |
| <i>Actitis macularia</i> | Spotted Sandpiper | S5B | | | | | CO | | |
| <i>Scolopax minor</i> | American Woodcock | S4B | | | | | PR | | |
| Ardeidae | Hérons & Bitterns | | | | | | | | |
| <i>Ardea herodias</i> | Great Blue Heron | S4 | | | | | CO | | |
| <i>Butorides virescens</i> | Green Heron | S4B | | | | | PO | | |
| Cathartidae | Vultures | | | | | | | | |
| <i>Cathartes aura</i> | Turkey Vulture | S5B,S3N | | | | | PR | | OB |
| Accipitridae | Hawks, Kites, Eagles & Allies | | | | | | | | |
| <i>Accipiter cooperii</i> | Cooper's Hawk | S4 | NAR | NAR | NS | No schedule | CO | | |
| <i>Accipiter striatus</i> | Sharp-shinned Hawk | S5 | NAR | NAR | NS | No schedule | CO | | |
| <i>Buteo jamaicensis</i> | Red-tailed Hawk | S5 | NAR | NAR | NS | No schedule | CO | | |
| Strigidae | Typical Owls | | | | | | | | |
| <i>Bubo virginianus</i> | Great Horned Owl | S4 | | | | | CO | | |
| <i>Megascops asio</i> | Eastern Screech-Owl | S4 | NAR | NAR | NS | No schedule | PR | | |
| Alcedinidae | Kingfishers | | | | | | | | |
| <i>Megaceryle alcyon</i> | Belted Kingfisher | S5B,S4N | | | | | CO | | |
| Picidae | Woodpeckers | | | | | | | | |
| <i>Colaptes auratus</i> | Northern Flicker | S5 | | | | | CO | | PR |
| <i>Dryobates pubescens</i> | Downy Woodpecker | S5 | | | | | CO | | PO |
| <i>Dryobates villosus</i> | Hairy Woodpecker | S5 | | | | | CO | | PR |
| <i>Dryocopus pileatus</i> | Pileated Woodpecker | S5 | | | | | CO | | |
| Falconidae | Caracaras & Falcons | | | | | | | | |
| <i>Falco sparverius</i> | American Kestrel | S4 | | | | | CO | | |
| Tyrannidae | Tyrant Flycatchers | | | | | | | | |
| <i>Contopus virens</i> | Eastern Wood-Pewee | S4B | SC | SC | SC | Schedule 1 | PR | X | |
| <i>Empidonax alnorum</i> | Alder Flycatcher | S5B | | | | | PO | | OB |
| <i>Empidonax minimus</i> | Least Flycatcher | S5B | | | | | PO | | |
| <i>Empidonax traillii</i> | Willow Flycatcher | S4B | | | | | PR | | |
| <i>Myiarchus crinitus</i> | Great Crested Flycatcher | S5B | | | | | CO | | |
| <i>Sayornis phoebe</i> | Eastern Phoebe | S5B | | | | | PO | | |
| <i>Tyrannus tyrannus</i> | Eastern Kingbird | S4B | | | | | CO | | |
| Vireonidae | Vireos | | | | | | | | |

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | OBBA* | NHIC Data** | NRSI Observed: Highest Level of Breeding Evidence |
|-----------------------------------|---|----------|-----------|---------------------------|---------------------------|---------------------------|-----------------|-------------|--|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | BSC et al. 2006 | MNR 2024 | NRSI Results from 2023 and 2025 |
| <i>Vireo gilvus</i> | Warbling Vireo | S5B | | | | | PR | | |
| <i>Vireo olivaceus</i> | Red-eyed Vireo | S5B | | | | | CO | | |
| Corvidae | Crows & Jays | | | | | | | | |
| <i>Corvus brachyrhynchos</i> | American Crow | S5 | | | | | CO | | PR |
| <i>Corvus corax</i> | Common Raven | S5 | | | | | | | OB |
| <i>Cyanocitta cristata</i> | Blue Jay | S5 | | | | | CO | | PR |
| Alaudidae | Larks | | | | | | | | |
| <i>Eremophila alpestris</i> | Horned Lark | S4 | | | | | PR | | |
| Hirundinidae | Swallows | | | | | | | | |
| <i>Hirundo rustica</i> | Barn Swallow | S4B | SC | SC | T | Schedule 1 | CO | | |
| <i>Petrochelidon pyrrhonota</i> | Cliff Swallow | S4S5B | | | | | CO | | |
| <i>Progne subis</i> | Purple Martin | S3B | | | | | PR | | |
| <i>Riparia riparia</i> | Bank Swallow | S4B | THR | T | T | Schedule 1 | PR | | |
| <i>Stelgidopteryx serripennis</i> | Northern Rough-winged Swallow | S4B | | | | | CO | | |
| <i>Tachycineta bicolor</i> | Tree Swallow | S4S5B | | | | | PR | | OB |
| Paridae | Chickadees & Titmice | | | | | | | | |
| <i>Baeolophus bicolor</i> | Tufted Titmouse | S3 | | | | | PR | X | |
| <i>Poecile atricapillus</i> | Black-capped Chickadee | S5 | | | | | CO | | PR |
| Sittidae | Nuthatches | | | | | | | | |
| <i>Sitta canadensis</i> | Red-breasted Nuthatch | S5 | | | | | CO | | |
| <i>Sitta carolinensis</i> | White-breasted Nuthatch | S5 | | | | | CO | | |
| Certhiidae | Creepers | | | | | | | | |
| <i>Certhia americana</i> | Brown Creeper | S5 | | | | | PO | | |
| Troglodytidae | Wrens | | | | | | | | |
| <i>Thryothorus ludovicianus</i> | Carolina Wren | S4 | | | | | CO | | PO |
| <i>Troglodytes aedon</i> | House Wren | S5B | | | | | CO | | PO |
| Polioptilidae | Gnatcatchers | | | | | | | | |
| <i>Polioptila caerulea</i> | Blue-gray Gnatcatcher | S4B | | | | | PR | | |
| Regulidae | Kinglets | | | | | | | | |
| <i>Corthylio calendula</i> | Ruby-crowned Kinglet | S5B, S3N | | | | | | | OB |
| Turdidae | Thrushes | | | | | | | | |
| <i>Catharus fuscescens</i> | Veery | S5B | | | | | PO | | |
| <i>Hylocichla mustelina</i> | Wood Thrush | S4B | SC | T | T | Schedule 1 | PR | | |
| <i>Turdus migratorius</i> | American Robin | S5 | | | | | CO | | CO |
| Mimidae | Mockingbirds, Thrashers & Allies | | | | | | | | |
| <i>Dumetella carolinensis</i> | Gray Catbird | S5B, S3N | | | | | CO | | |
| <i>Mimus polyglottos</i> | Northern Mockingbird | S4 | | | | | CO | | |
| <i>Toxostoma rufum</i> | Brown Thrasher | S4B | | | | | CO | | OB |
| Sturnidae | Starlings | | | | | | | | |
| <i>Sturnus vulgaris</i> | European Starling | SNA | | | | | CO | | |
| Bombycillidae | Waxwings | | | | | | | | |
| <i>Bombycilla cedrorum</i> | Cedar Waxwing | S5 | | | | | CO | | OB |
| Passeridae | Old World Sparrows | | | | | | | | |
| <i>Passer domesticus</i> | House Sparrow | SNA | | | | | CO | | PR |
| Fringillidae | Finches & Allies | | | | | | | | |
| <i>Haemorhous mexicanus</i> | House Finch | SNA | | | | | CO | | |
| <i>Spinus tristis</i> | American Goldfinch | S5 | | | | | CO | | PR |
| Emberizidae | New World Sparrows & Allies | | | | | | | | |
| <i>Ammodramus savannarum</i> | Grasshopper Sparrow | S4B | SC | SC | SC | Schedule 1 | PO | | |
| <i>Junco hyemalis</i> | Dark-eyed Junco | S5 | | | | | | | OB |
| <i>Melospiza melodia</i> | Song Sparrow | S5 | | | | | CO | | PO |
| <i>Passerculus sandwichensis</i> | Savannah Sparrow | S5B, S3N | | | | | CO | | |
| <i>Pipilo erythrophthalmus</i> | Eastern Towhee | S4B, S3N | | | | | PR | | |
| <i>Poocetes gramineus</i> | Vesper Sparrow | S4B | | | | | PO | | |
| <i>Spizella passerina</i> | Chipping Sparrow | S5B, S3N | | | | | CO | | |
| <i>Spizella pusilla</i> | Field Sparrow | S4B, S3N | | | | | PO | | OB |
| Icteridae | Troupials & Allies | | | | | | | | |
| <i>Agelaius phoeniceus</i> | Red-winged Blackbird | S5 | | | | | CO | | OB |

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | OBBA* | NHIC Data** | NRSI Observed: Highest Level of Breeding Evidence |
|--------------------------------|--|----------|-----------|---------------------------|---------------------------|---------------------------|-----------------|-------------|--|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | BSC et al. 2006 | MNR 2024 | NRSI Results from 2023 and 2025 |
| <i>Dolichonyx oryzivorus</i> | Bobolink | S4B | THR | SC | T | Schedule 1 | PO | | |
| <i>Icterus galbula</i> | Baltimore Oriole | S4B | | | | | CO | | |
| <i>Molothrus ater</i> | Brown-headed Cowbird | S5 | | | | | CO | | |
| <i>Quiscalus quiscula</i> | Common Grackle | S5 | | | | | CO | | PR |
| <i>Sturnella magna</i> | Eastern Meadowlark | S4B, S3N | THR | T | T | Schedule 1 | PR | | |
| Parulidae | Wood Warblers | | | | | | | | |
| <i>Geothlypis philadelphia</i> | Mourning Warbler | S5B | | | | | PO | | |
| <i>Geothlypis trichas</i> | Common Yellowthroat | S5B, S3N | | | | | PR | | |
| <i>Seiurus aurocapilla</i> | Ovenbird | S5B | | | | | PO | | |
| <i>Setophaga coronata</i> | Yellow-rumped Warbler | S5B, S4N | | | | | | | OB |
| <i>Setophaga pensylvanica</i> | Chestnut-sided Warbler | S5B | | | | | PO | | |
| <i>Setophaga petechia</i> | Yellow Warbler | S5B | | | | | CO | | |
| <i>Setophaga pinus</i> | Pine Warbler | S5B, S3N | | | | | PR | | |
| <i>Setophaga ruticilla</i> | American Redstart | S5B | | | | | PR | | |
| Cardinalidae | Cardinals, Grosbeaks & Allies | | | | | | | | |
| <i>Cardinalis cardinalis</i> | Northern Cardinal | S5 | | | | | CO | | CO |
| <i>Passerina cyanea</i> | Indigo Bunting | S5B | | | | | CO | | |
| <i>Pheucticus ludovicianus</i> | Rose-breasted Grosbeak | S5B | | | | | CO | | |
| <i>Piranga olivacea</i> | Scarlet Tanager | S5B | | | | | PO | | |
| Total | | | | | | | 91 | 4 | 27 |

*OBBA Atlas Square: 17PJ01

**NHIC Atlas Square: 17PJ0413

References

Bird Studies Canada (BSC), Environment Canada's Canadian Wildlife Service, Ontario Nature, Ontario Field Ornithologists and Ontario Ministry of Natural Resources. 2006. Ontario Breeding Bird Atlas Database, 31 January 2008. <https://www.birdsontario.org/jsp/datasummaries.jsp>

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>

Ministry of Natural Resources (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Appendix V
Herpetofauna Species Reported from the Study Area

Reptile and Amphibian Species Reported from the Study Area - 1493 Sixth Line, Oakville (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | ORAA* | NHIC Data** | NRSI Observed |
|---|--|----------|-----------|---------------------------|---------------------------|---------------------------|---------------------|-------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Ontario Nature 2019 | MNR 2024 | NRSI Results from 2023 and 2025 |
| Turtles | | | | | | | | | |
| <i>Chelydra serpentina</i> | Snapping Turtle | S4 | SC | SC | SC | Schedule 1 | X | | |
| <i>Chrysemys picta marginata</i> | Midland Painted Turtle | S4 | | SC | SC | Schedule 1 | X | | |
| <i>Graptemys geographica</i> | Northern Map Turtle | S3 | SC | SC | SC | Schedule 1 | X | | |
| <i>Trachemys scripta</i> | Pond Slider | SNA | | | | | X | | |
| Snakes | | | | | | | | | |
| <i>Diadophis punctatus</i> | Northern Ring-necked Snake | S4 | | | | | X | | |
| <i>Lampropeltis triangulum</i> | Eastern Milksnake | S4 | NAR | SC | SC | Schedule 1 | X | | |
| <i>Opheodrys vernalis</i> | Smooth Greensnake | S4 | | | | | X | | |
| <i>Nerodia sipedon sipedon</i> | Northern Watersnake | S5 | NAR | NAR | NS | No schedule | X | | |
| <i>Storeria dekayi</i> | Dekay's Brownsnake | S5 | NAR | NAR | NS | No schedule | X | | |
| <i>Storeria occipitomaculata</i> | Red-bellied Snake | S5 | | | | | X | | |
| <i>Thamnophis sirtalis sirtalis</i> | Eastern Gartersnake | S5 | | | | | X | | |
| Salamanders | | | | | | | | | |
| <i>Ambystoma jeffersonianum</i> | Jefferson Salamander | S2 | END | E | E | Schedule 1 | X | | |
| <i>Ambystoma laterale</i> - (2) <i>jeffersonianum</i> | Unisexual Ambystoma (Jefferson Salamander) | S2 | END | E | E | Schedule 1 | X | | |
| <i>Ambystoma maculatum</i> | Spotted Salamander | S4 | | | | | X | | |
| <i>Notophthalmus viridescens viridescens</i> | Red-spotted Newt | S5 | | | | | X | | |
| <i>Plethodon cinereus</i> | Eastern Red-backed Salamander | S5 | | | | | X | | |
| Frogs and Toads | | | | | | | | | |
| <i>Anaxyrus americanus</i> | American Toad | S5 | | | | | X | | |
| <i>Dryophytes versicolor</i> | Gray Treefrog | S5 | | | | | X | | |
| <i>Pseudacris triseriata</i> pop. 2 | Western Chorus Frog (Great Lakes / St. L.) | S4 | NAR | T | T | Schedule 1 | X | | |
| <i>Pseudacris crucifer</i> | Spring Peeper | S5 | | | | | X | | |
| <i>Lithobates clamitans</i> | Green Frog | S5 | | | | | X | | |
| <i>Lithobates pipiens</i> | Northern Leopard Frog | S5 | NAR | NAR | NS | No schedule | X | | |
| <i>Lithobates sylvaticus</i> | Wood Frog | S5 | | | | | X | | |
| Total | | | | | | | 23 | 0 | 0 |

*ORAA Atlas Square: 17PJ01

**NHIC Atlas Square: 17PJ0413

References

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>

Ministry of Natural Resources (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Ontario Nature. 2019. Ontario Reptile and Amphibian Atlas Program: Interactive Range Maps. Accessed October 2019.

Appendix VI
Mammal Species Reported from the Study Area

Mammal Species Reported from the Study Area - 1493 Sixth Line, Oakville (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Ontario Mammal Atlas | NHIC Data** | NRSI Observed |
|----------------------------------|---|----------|-----------|---------------------------|---------------------------|---------------------------|----------------------|-------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Dobbyn 1994 | MNR 2024 | NRSI Results from 2023 and 2025 |
| Didelphimorphia | Opossums | | | | | | | | |
| <i>Didelphis virginiana</i> | Virginia Opossum | S4 | | | | | X | | |
| Eulipotyphla | Shrews, Moles, Hedgehogs, and Allies | | | | | | | | |
| <i>Blarina brevicauda</i> | Northern Short-tailed Shrew | S5 | | | | | X | | |
| <i>Condylura cristata</i> | Star-nosed Mole | S5 | | | | | X | | |
| <i>Parascalops breweri</i> | Hairy-tailed Mole | S4 | | | | | X | | |
| <i>Sorex cinereus</i> | Masked Shrew | S5 | | | | | X | | |
| <i>Sorex fumeus</i> | Smoky Shrew | S5 | | | | | X | | |
| <i>Sorex hoyi</i> | Pygmy Shrew | S4 | | | | | X | | |
| <i>Sorex palustris</i> | Water Shrew | S5 | | | | | X | | |
| Chiroptera | Bats | | | | | | | | |
| <i>Eptesicus fuscus</i> | Big Brown Bat | S4 | | | | | X | | |
| <i>Lasionycteris noctivagans</i> | Silver-haired Bat | S3 | END | E | NS | No schedule | X | | |
| <i>Lasius borealis</i> | Eastern Red Bat | S3 | END | E | NS | No schedule | X | | |
| <i>Lasius cinereus</i> | Hoary Bat | S3 | END | E | NS | No schedule | X | | |
| <i>Myotis lucifugus</i> | Little Brown Myotis | S3 | END | E | E | Schedule 1 | X | | |
| <i>Myotis septentrionalis</i> | Northern Myotis | S3 | END | E | E | Schedule 1 | X | | |
| <i>Perimyotis subflavus</i> | Tri-colored Bat | S3? | END | E | E | Schedule 1 | X | | |
| Lagomorpha | Rabbits and Hares | | | | | | | | |
| <i>Lepus americanus</i> | Snowshoe Hare | S5 | | | | | X | | |
| <i>Lepus europaeus</i> | European Hare | SNA | | | | | X | | |
| <i>Sylvilagus floridanus</i> | Eastern Cottontail | S5 | | | | | X | | X |
| Rodentia | Rodents | | | | | | | | |
| <i>Castor canadensis</i> | Beaver | S5 | | | | | X | | |
| <i>Erethizon dorsatum</i> | Porcupine | S5 | | | | | X | | |
| <i>Glaucomys sabrinus</i> | Northern Flying Squirrel | S5 | | | | | X | | |
| <i>Marmota monax</i> | Woodchuck | S5 | | | | | X | | |
| <i>Microtus pennsylvanicus</i> | Meadow Vole | S5 | | | | | X | | |
| <i>Mus musculus</i> | House Mouse | SNA | | | | | X | | |
| <i>Ondatra zibethicus</i> | Muskrat | S4 | | | | | X | | |
| <i>Peromyscus leucopus</i> | White-footed Mouse | S5 | | | | | X | | |
| <i>Peromyscus maniculatus</i> | Deer Mouse | S5 | | | | | X | | |
| <i>Rattus norvegicus</i> | Norway Rat | SNA | | | | | X | | |
| <i>Sciurus carolinensis</i> | Eastern Gray Squirrel | S5 | | | | | X | | X |
| <i>Tamias striatus</i> | Eastern Chipmunk | S5 | | | | | X | | X |
| <i>Tamiasciurus hudsonicus</i> | Red Squirrel | S5 | | | | | X | | |
| <i>Zapus hudsonius</i> | Meadow Jumping Mouse | S5 | | | | | X | | |
| Canidae | Canines | | | | | | | | |
| <i>Canis latrans</i> | Coyote | S5 | | | | | X | | |
| <i>Vulpes vulpes</i> | Red Fox | S5 | | | | | X | | |
| Mephitidae | Skunks and Stink Badgers | | | | | | | | |
| <i>Mephitis mephitis</i> | Striped Skunk | S5 | | | | | X | | |
| Mustelidae | Weasels and Allies | | | | | | | | |
| <i>Lontra canadensis</i> | North American River Otter | S5 | | | | | X | | |
| <i>Mustela richardsonii</i> | American Ermine | S5 | | | | | X | | |
| <i>Neogale frenata</i> | Long-tailed Weasel | S4 | | | | | X | | |
| <i>Neogale vison</i> | American Mink | S4 | | | | | X | | |
| Procyonidae | Raccoons and Allies | | | | | | | | |
| <i>Procyon lotor</i> | Northern Raccoon | S5 | | | | | X | | X |
| Ursidae | Bears | | | | | | | | |
| <i>Ursus americanus</i> | American Black Bear | S5 | NAR | NAR | NS | No schedule | X | | |
| Artiodactyla | Deer and Bison | | | | | | | | |
| <i>Alces alces</i> | Moose | S5 | | | | | X | | |
| <i>Odocoileus virginianus</i> | White-tailed Deer | S5 | | | | | X | | |
| Total | | | | | | | 43 | 0 | 4 |

*Mammal Atlas Square Number: PU

**NHIC Atlas Square: 17PJ0413

References

Dobbyn, J.S. 1994. Atlas of the Mammals of Ontario. Don Mills, Federation of Ontario Naturalists. 120p.

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>

Ministry of Natural Resources (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Appendix VII
Lepidoptera Species Reported from the Study Area

Butterfly Species Reported from the Study Area - 1493 Sixth Line, Oakville EIS (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Ontario Butterfly Atlas* | NHIC Data** | NRSI Observed |
|------------------------------------|--|----------|-----------|---------------------------|---------------------------|---------------------------|--------------------------|-------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Macnaughton et al. 2025 | MNR 2024 | NRSI Results from 2023 and 2025 |
| Hesperiidae | Skippers | | | | | | | | |
| <i>Anatrytone logan</i> | Delaware Skipper | S4 | | | | | X | | |
| <i>Ancyloxypha numitor</i> | Least Skipper | S5 | | | | | X | | |
| <i>Epgeryx clarus</i> | Silver-spotted Skipper | S4 | | | | | X | | |
| <i>Erynnis baptisiae</i> | Wild Indigo Duskywing | S4 | | | | | X | | |
| <i>Erynnis icelus</i> | Dreamy Duskywing | S5 | | | | | X | | |
| <i>Erynnis juvenalis</i> | Juvenal's Duskywing | S5 | | | | | X | | |
| <i>Euphyes dion</i> | Dion Skipper | S4 | | | | | X | | |
| <i>Euphyes vestris</i> | Dun Skipper | S5 | | | | | X | | |
| <i>Hylephila phyleus</i> | Fiery Skipper | SNA | | | | | X | | |
| <i>Pholisora catullus</i> | Common Sootywing | S4 | | | | | X | | |
| <i>Polites mystic</i> | Long Dash Skipper | S5 | | | | | X | | |
| <i>Polites origenes</i> | Crossline Skipper | S4 | | | | | X | | |
| <i>Polites themistocles</i> | Tawny-edged Skipper | S5 | | | | | X | | |
| <i>Thorybes pylades</i> | Northern Cloudywing | S5 | | | | | X | | |
| <i>Thymelicus lineola</i> | European Skipper | SNA | | | | | X | | |
| Papilionidae | Swallowtails | | | | | | | | |
| <i>Battus philenor</i> | Pipevine Swallowtail | SNA | | | | | X | | |
| <i>Heraclides cresphontes</i> | Giant Swallowtail | S4 | | | | | X | | |
| <i>Papilio glaucus</i> | Eastern Tiger Swallowtail | S5 | | | | | | | X |
| <i>Papilio polyxenes</i> | Black Swallowtail | S5 | | | | | X | | X |
| Pieridae | Whites and Sulphurs | | | | | | | | |
| <i>Colias eurytheme</i> | Orange Sulphur | S5 | | | | | X | | |
| <i>Colias philodice</i> | Clouded Sulphur | S5 | | | | | X | | X |
| <i>Pieris oleracea</i> | Mustard White | S4 | | | | | X | | |
| <i>Pieris rapae</i> | Cabbage White | SNA | | | | | X | | X |
| <i>Pieris virginensis</i> | West Virginia White | S3 | SC | | | | X | | |
| Lycaenidae | Harvesters, Coppers, Hairstreaks, Blues | | | | | | | | |
| <i>Callophrys niphon</i> | Eastern Pine Elfin | S5 | | | | | X | | |
| <i>Celastrina lucia</i> | Northern Spring Azure | S5 | | | | | X | | |
| <i>Celastrina neglecta</i> | Summer Azure | S5 | | | | | X | | |
| <i>Celastrina sp.</i> | Azure species | SNA | | | | | X | | |
| <i>Cupido comyntas</i> | Eastern Tailed Blue | S5 | | | | | X | | |
| <i>Glaucopsyche lygdamus</i> | Silvery Blue | S5 | | | | | X | | |
| <i>Polyommatus icarus</i> | European Common Blue | SNA | | | | | X | | |
| <i>Satyrium acadica</i> | Acadian Hairstreak | S4 | | | | | X | | |
| <i>Satyrium calanus</i> | Banded Hairstreak | S4 | | | | | X | | |
| <i>Satyrium caryaevorus</i> | Hickory Hairstreak | S4 | | | | | X | | |
| <i>Satyrium liparops</i> | Striped Hairstreak | S5 | | | | | X | | |
| Nymphalidae | Brush-footed Butterflies | | | | | | | | |
| <i>Aglais milberti</i> | Milbert's Tortoiseshell | S5 | | | | | X | | |
| <i>Boloria bellona</i> | Meadow Fritillary | S5 | | | | | X | | |
| <i>Cercyonis pegala</i> | Common Wood-Nymph | S5 | | | | | X | | |
| <i>Chlosyne nycteis</i> | Silvery Checkerspot | S5 | | | | | X | | |
| <i>Coenonympha californica</i> | Common Ringlet | S5 | | | | | X | | |
| <i>Danaus plexippus</i> | Monarch | S2N,S4B | SC | E | E | Schedule 1 | X | | |
| <i>Euphydryas phaeton</i> | Baltimore Checkerspot | S4 | | | | | X | | |
| <i>Euptoieta claudia</i> | Variegated Fritillary | SNA | | | | | X | | |
| <i>Junonia coenia</i> | Common Buckeye | SNA | | | | | X | | |
| <i>Lethe anhedon</i> | Northern Pearly-Eye | S5 | | | | | X | | |
| <i>Lethe appalachia</i> | Appalachian Brown | S4 | | | | | X | | |
| <i>Lethe eurydice</i> | Eyed Brown | S5 | | | | | X | | |
| <i>Libytheana carinenta</i> | American Snout | SNA | | | | | X | | |
| <i>Limenitis archippus</i> | Viceroy | S5 | | | | | X | | |
| <i>Limenitis arthemis arthemis</i> | White Admiral | S5 | | | | | X | | |
| <i>Limenitis arthemis astyanax</i> | Red-spotted Purple | S5 | | | | | X | | |
| <i>Megisto cymela</i> | Little Wood-Satyr | S5 | | | | | X | | |
| <i>Nymphalis antiopa</i> | Mourning Cloak | S5 | | | | | X | | |
| <i>Nymphalis l-album</i> | Compton Tortoiseshell | S5 | | | | | X | | |
| <i>Phyciodes cocyta</i> | Northern Crescent | S5 | | | | | X | | |
| <i>Phyciodes tharos</i> | Pearl Crescent | S4 | | | | | X | | |
| <i>Polygonia comma</i> | Eastern Comma | S5 | | | | | X | | |

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Ontario Butterfly Atlas* | NHIC Data** | NRSI Observed |
|----------------------------------|---------------|----------|-----------|---------------------------|---------------------------|---------------------------|--------------------------|-------------|---------------------------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | Macnaughton et al. 2025 | MNR 2024 | NRSI Results from 2023 and 2025 |
| <i>Polygonia interrogationis</i> | Question Mark | S5 | | | | | X | | |
| <i>Polygonia progne</i> | Gray Comma | S5 | | | | | X | | |
| <i>Vanessa atalanta</i> | Red Admiral | S5B | | | | | X | | |
| <i>Vanessa cardui</i> | Painted Lady | S5B | | | | | X | | |
| <i>Vanessa virginiensis</i> | American Lady | S5 | | | | | X | | |
| Total | | | | | | | 61 | 0 | 4 |

*TEA Atlas Square: 17PJ01

**NHIC Atlas Square: 17PJ0413

References

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Macnaughton A., Layberry R., Cavašin R., Edwards B., and C. Jones. 2025. Ontario Butterfly Atlas. Updated February 2025. Available: <https://www.ontarioinsects.org/atlas/index.html>

Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>

Ministry of Natural Resources (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Appendix VIII
Odonata Species Reported from the Study Area

Odonate Species Reported from the Study Area - 1493 Sixth Line, Oakville (Project #3096A)

| Scientific Name | Common Name | SRANK | SARO | COSEWIC | SARA | SARA Schedule | Odonate Atlas* | NHIC Data** |
|----------------------------------|----------------------------------|----------|-----------|---------------------------|---------------------------|---------------------------|----------------|-------------|
| | | MNR 2025 | MECP 2024 | Government of Canada 2024 | Government of Canada 2024 | Government of Canada 2024 | OOAD 2023 | MNR 2024 |
| Calopterygidae | Broadwinged Damselflies | | | | | | | |
| <i>Calopteryx aequabilis</i> | River Jewelwing | S5 | | | | | X | |
| <i>Calopteryx maculata</i> | Ebony Jewelwing | S5 | | | | | X | |
| <i>Heterina americana</i> | American Rubyspot | S4 | | | | | X | |
| Lestidae | Spreadwings | | | | | | | |
| <i>Lestes dryas</i> | Emerald Spreadwing | S5 | | | | | X | |
| Coenagrionidae | Narrow-winged Damselflies | | | | | | | |
| <i>Argia fumipennis violacea</i> | Violet Dancer | S5 | | | | | X | |
| <i>Argia moesta</i> | Powdered Dancer | S5 | | | | | X | |
| <i>Enallagma antennatum</i> | Rainbow Bluet | S4 | | | | | X | |
| <i>Enallagma civile</i> | Familiar Bluet | S5 | | | | | X | |
| <i>Enallagma exulans</i> | Stream Bluet | S5 | | | | | X | |
| <i>Ichnura posita</i> | Fragile Forktail | S4 | | | | | X | |
| <i>Ichnura verticalis</i> | Eastern Forktail | S5 | | | | | X | |
| Aeshnidae | Darners | | | | | | | |
| <i>Aeshna canadensis</i> | Canada Darter | S5 | | | | | X | |
| <i>Anax junius</i> | Common Green Darter | S5 | | | | | X | |
| <i>Basiaeschna janata</i> | Springtime Darter | S5 | | | | | X | |
| Gomphidae | Clubtails | | | | | | | |
| <i>Ophiogomphus rupisulensis</i> | Rusty Snaketail | S4 | | | | | X | |
| <i>Phanogomphus lividus</i> | Ashy Clubtail | S4 | | | | | X | |
| Libellulidae | Skimmers | | | | | | | |
| <i>Erythemis simplicicollis</i> | Eastern Pondhawk | S5 | | | | | X | |
| <i>Libellula luctuosa</i> | Widow Skimmer | S5 | | | | | X | |
| <i>Libellula pulchella</i> | Twelve-spotted Skimmer | S5 | | | | | X | |
| <i>Libellula quadrimaculata</i> | Four-spotted Skimmer | S5 | | | | | X | |
| <i>Libellula semifasciata</i> | Painted Skimmer | S3 | | | | | X | |
| <i>Pachydiplax longipennis</i> | Blue Dasher | S5 | | | | | X | |
| <i>Pantala hymenaea</i> | Spot-winged Glider | S4 | | | | | X | |
| <i>Plathemis lydia</i> | Common Whitetail | S5 | | | | | X | |
| <i>Tamea lacerata</i> | Black Saddlebags | S4 | | | | | X | |
| <i>Tamea onusta</i> | Red Saddlebags | SNA | | | | | X | |
| Total | | | | | | | 26 | 0 |

*Odonate Atlas Square Numbers: 17PJ01

**NHIC Atlas Square: 17PJ0413

References

Government of Canada. 2023. Species at Risk Public Registry: Species Search. COSEWIC Last Assessment Date: 2023-05-05. Available: <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>
Ministry of the Environment, Conservation, and Parks (MECP). 2024. Species at Risk in Ontario. Published: 2018-07-12. Updated: 2024-04-19. Available: <https://www.ontario.ca/page/species-risk-ontario>
Ministry of Natural Resources (MNR). 2025. Natural Heritage Information Centre (NHIC): Species List for Ontario. Published: 2014-07-17. All Species List Updated: 2025-03-30. Available: <https://www.ontario.ca/page/get-natural-heritage-information>
Ministry of Natural Resources (MNR). 2024. Natural Heritage Information Centre (NHIC): Make a Natural Heritage Area Map Application. Published: 2014-07-17. Updated 2024-10-30. Available: <https://www.ontario.ca/page/make-natural-heritage-area-map>
Ontario Odonata Atlas Database (OOAD). 2023. Natural Heritage Information Centre, Ontario Ministry of Natural Resources and Forestry. Species data by 10x10 km square accessed on June 8, 2023